

Valencia SWCD Project Tome 11.09

10-year Monitoring Report

2022



Prepared by

Alex Makowicki and Patrick Clay Goetsch

With the 2011/2012 work of Joe Zebrowski,

and the 2016 work of Kathryn R Mahan & Daniel Hernandez

New Mexico Forest and watershed Restoration Institute

For the Greater Rio Grande Watershed Alliance



Contents

Acronyms and Abbreviations	3
Purpose of Report	4
Ecological Context of Bosque Restoration	4
Monitoring and Field Methods	5
Original (2012) protocols	5
5 and 10-year revisits (2016 and 2022) protocols	6
Personnel Involved.....	7
Bosque Ecological Monitoring Program Sites.....	8
Tome Project.....	8
Tome Site Summary	10
Tome 2012 & 2022 - Observed plant species summary	11
Tree Component	13
Understory Component	14
Next steps (monitoring)	19
References	19
Appendix I – Plot Coordinates Table.....	20
Appendix II - Modified Hink and Ohmart categories, from NMRAM.....	21
Appendix III – Sample Datasheet	24
Appendix IV – Fuels Transect Data Sheet	28
Appendix V – Photo Pages	29

Acronyms and Abbreviations

Acronym, Abbreviation, or Term	Explanation or Definition as used by NMFWR
FSA	Farm Service Agency, a department of the USDA
GIS	Geographic Information Systems
GRGWA	Greater Rio Grande Watershed Alliance
LIDAR	Light detecting and ranging, a remote sensing technique using light to gather elevation data
NHNM	Natural Heritage New Mexico
NMDGF	New Mexico Department of Game and Fish
NMED SWQB	New Mexico Environment Department Surface Water Quality Bureau
NMFWR	New Mexico Forest and Watershed Restoration Institute
NMHU	New Mexico Highlands University
NMRAM	New Mexico Rapid Assessment Method, version 2.1
NRCS	Natural Resource Conservation Service
PC	Plot center
RGIS	Resource Geographic Information System
SWCD	Soil and Water Conservation District
USDA	United States Department of Agriculture
USGS	United States Geological Survey
WQCC	Water Quality Control Commission
WSS	Web Soil Survey, a soils database of the NRCS

Purpose of Report

This report covers pre-treatment and 5 and 10-year-post-treatment vegetation monitoring assessments performed on non-native phreatophyte removal projects near Belen, NM submitted by the Valencia Soil and Water Conservation District to the Greater Rio Grande Watershed Alliance in 2011. Following a discussion of the ecological context, and our monitoring methods, we present pertinent background, observations, and assessment results for each project.

Ecological Context of Bosque Restoration

Neither the challenges nor the importance of working in the bosque and other riparian areas in New Mexico today should be underestimated. According to the New Mexico Department of Game and Fish Conservation Division, wetlands and riparian areas comprise approximately 0.6 percent of all land in New Mexico (2012). Despite this small percentage, estimates of New Mexican vertebrate species depending on wetland and riparian habitat for their survival ranges from 55% (New Mexico Department of Game and Fish Conservation Services Division, 2012) to 80% (Audubon New Mexico, 2013). These areas also provide flood mitigation, filtration of sediment and pollutants, and water for a variety of purposes including groundwater recharge (Audubon New Mexico, 2013). In addition, native vegetation such as cottonwoods have cultural significance to many communities.

As much as these areas are disproportionately important to ecosystems and human communities, they are equally disproportionately impacted by disturbance. Anthropogenic impacts with major consequences for our riparian areas include dams, reservoirs, levees, channelization, acequias and ditches, jetty jacks, riprap and Gabion baskets, urbanization, removal of native phreatophytes, grazing by domestic livestock, excessive grazing pressure by native ungulate populations absent natural predation cycles, beaver removal, logging, mining, recreation, transportation, introduction and spread of invasive exotic species, groundwater extraction, altered fire and flood regimes drought and climate change (Committee on Riparian Zone Functioning and Strategies for Management, et al., 2002). Statewide, it is estimated that as much as 90% of New Mexico's historical riparian areas have been lost (Audubon New Mexico, 2013), and approximately 39% of our remaining perennial stream miles are impaired (New Mexico Department of Game and Fish Conservation Services Division, 2012).

New Mexico is fortunate enough to have the Middle Rio Grande Bosque, the largest remaining bosque in the Southwest (USDA USFS, 1996). However, over the past two decades, the number of fires in the bosque has been increasing. Historically, the primary disturbance regime in the bosque has been flooding, not fire, which means the system is not fire-adapted. In fact, native species like cottonwood resprout from their roots after floods and need wet soils to germinate from seed. Flooding also promotes decomposition of organic material and keeps the soil moist which reduces the likelihood of fire. Today, overbank flow is uncommon in many areas of the Rio Grande due to the heavy alteration of the channel and flow regimes (two obvious examples are the structures defining the upper and lower extent of the Middle Rio Grande: Cochiti Dam and Elephant Butte Reservoir). This has led to low fuel moisture content and high fuel loads, as well as increased human presence in the riparian area. As a result, bosque fires are more common and more severe: they kill cottonwoods and other native species, creating spaces which are filled by non-native species such as salt cedar, Russian olive, Siberian elm, and Tree-of-Heaven. We are constantly learning more about how these species can exploit and encourage a riparian fire regime, in addition to many other changes they bring to ecosystems.

Efforts geared toward the removal of these nonnative species can help to reduce fire risk, preserve native vegetation, and be part of a larger effort to restore the bosque and the watershed as a whole to a more natural and functional ecosystem. The Greater Rio Grande Watershed Alliance (GRGWA) has been working on these issues with a variety of collaborating organizations and agencies within the Rio Grande basin for several years. Since 2013, the New Mexico Forest and Watershed Restoration Institute (NMFWRRI) has been working with GRGWA and the Claunch-Pinto Soil and Water Conservation District (SWCD) to begin construction of a geodatabase for all of GRGWA's non-native phreatophyte removal projects as well as to perform the formal pre- and post-treatment monitoring, utilizing a range of field methods as well as LIDAR analysis where appropriate and available.

Monitoring and Field Methods

Original (2012) protocols

Due to the short timeframe between project selection and implementation in 2011/2012, only a narrow window was available to perform pre-treatment monitoring. That window was outside the optimum season for performing vegetation monitoring in this type of landscape. For that reason, a hasty monitoring protocol was developed. This protocol was based on placing photo point plots at locations distributed across the project area and representative of the diversity of the project area. In addition, an estimate of ground and canopy cover by percent within a 1/10-acre circular plot centered at the photo point was determined using ocular estimates. Overstory canopy was determined for a 1/10-acre circular area, also centered at the photo point. Finally, a Hink & Ohmart style vegetation structure assessment was performed. Vegetation species that were observed at each plot and in the project area were recorded. The plot size and density of observations limit the utility of this monitoring for describing overall site conditions or for generating any meaningful statistics.

Cover (%)											
Tree canopy	Seedlings/saplings <5' / 5 – 15'		Shrubs		Gramanoid	Forbs	Litter	Bare Soil	Rock	Gravel	Water or wet

Figure 1. Categories used for 2012 percent cover estimates.

A base map of the project location was constructed using project boundary data provided by New Mexico State Forestry. Planned photo points were selected by visual inspection of May 2011 true-color digital orthorectified aerial photography obtained from the United States Department of Agriculture (<http://datagateway.nrcs.usda.gov/>). A GIS file for the photo point plots was created using ArcGIS software. Coordinates were derived from the GIS file and loaded into a Garmin GPS 60 CSx Global Positioning System and a Trimble 2005 GeoXM Global Positioning System. The Garmin GPS was used to navigate to the general location of the planned photo point. The actual location of the photo point was determined by visual inspection of the area and selection was based on the ability to physically occupy a position at or near the planned point. The coordinates of the photo point were then collected using the more precise Trimble GeoXM GPS.

Once the plot location was determined, a 1/100-acre radius plot was established by placing pin-flags at 11' 9" from plot center in each cardinal direction. Photos were taken from plot center in each cardinal direction and from a distance north of plot center (66', where possible) toward plot center. Ocular estimates were made of understory canopy and ground cover within the 1/100 plot. Overstory canopy cover was estimated using a concave spherical densiometer, with measurements made in four cardinal directions, approximately mid-way between plot center and the edge of the 1/100-acre plot. This method provides an estimate of canopy cover for a 1/10-acre area centered on the plot. A Hink & Ohmart structure class determination was made using a worksheet developed by SWCA Environmental Consultants (see datasheet example in Appendix III). Finally, plant species observed within the 1/10 area around the plot were recorded, as were other comments document conditions at the plot.




Figure 2. example of plot layout. The outer circle represents the 1/10 acre plot and the blue circle is the 1/100 plot

5 and 10-year revisits (2016 and 2022) protocols

To allow comparisons between site conditions, the original site protocols were employed for the 5 and 10-year revisits as well as newer protocols for the 10-year revisit.

Plot locations as recorded in 2011 and 2016 were found using a Garmin GPS, and all plot setup and measurements were the same as in 2011 and 2016, with a few exceptions. In 2016 a ground cover category was added for plant basal/bole, which was omitted from the ground cover in 2011. Further, for both 2016 and 2022 monitoring, in addition to the original Hink and Ohmart structural classification, we recorded the structure type within a modified Hink and Ohmart classification system (see Appendix II). This second Hink and Ohmart-based system is used by the NMED as part of the modified NMRAM protocol employed for pre-treatment monitoring on GRGWA projects beginning in 2013. Additions in 2022 were the inclusion of NMFWR's Riparian Common Stand Exam-based protocols (https://nmfwri.org/wpcontent/uploads/2020/07/GRGWA_plotprotocols_Instructions_datasheets_with_cheatsheets_3.1.2020km.pdf) which added measurements of soil texture; ground and aerial cover on the entire plot as well as aerial cover by individual species, seedling and sapling tallies and individual tree measurements (Appendix III). Individual tree measurements included establishing a witness tree when available, measuring tree height, diameter at breast height (DBH), live crown base height and overall health of the tree. Fuel transects were also established. (Appendix IV).

For the sake of continuity, site visits were made around the same time of year as 5 and 10 years prior, even though this was not the ideal season for plant identification in either case. It is worth noting that

the winter of 2016/2017 was warmer than the winter of 2011/2012, so even though site visits were conducted around the same time of year, plant communities differed. This is especially obvious in the photographs (Appendix V).




Figure 3. Example of fuels transect

Personnel Involved

2012 Monitoring Team:

- Joe Zebrowski, New Mexico Forest and Watershed Restoration Institute
- Jill Wick, New Mexico Department of Game and Fish (Sites B1 and B2)
- Dave Lightfoot, SWCA Environmental Consultants (Sites B3 and B4)
- Cody Stropki, SWCA Environmental Consultants (Sites B3 and B4)

2016 Monitoring Team:

- Kathryn R Mahan, Ecological Monitoring Specialist
- Christopher B Martinez, Monitoring Technician (NMHU Student Intern)
- Daniel Hernandez, Ecological Monitoring Technician

2022 Monitoring Team:

- Alex Makowicki, Monitoring technician
- Clay Goetsch, Monitoring technician
- Jordan Martinez, Monitoring technician
-

Other persons contacted 2022:

- Yasmeen Najmi, Middle Rio Grande Conservancy District

Bosque Ecological Monitoring Program Sites

Three Bosque Ecological Monitoring Program (BEMP) monitoring sites were randomly selected using ArcMap 10.4.1. Sites were labeled, T_1, T_2, and T_3 (Fig.1). These sites were likely disturbed during the treatment activity. GRGWA monitoring now strives to integrate BEMP monitoring into the overall project monitoring scheme.



Figure 4. Tome plot locations. 11.09_1, 11.09_2 and 11.09_3.

Tome Project

The project is located within Valencia County, NM, west of the village of Tome. It is on the east side of the Rio Grande, between the levee road and river (Fig.4).

The nearby city of Los Lunas receives an average of 9.75 inches of precipitation annually. The average high temperature is 94 degrees in July, and the average low is 18 in December and January (U.S. Climate Data, 2017). According to the NRCS Web Soil Survey, the project area is comprised of <1% Riverwash

and the remainder Mixed alluvial land. Ecological sites within this project include R042XA055NM Salty Bottomland (USDA NRCS, 2016).

Salty Bottomland can support a range of plant communities which typically include cottonwood, salt cedar, mixed exotics (dominated by Russian olive/ Russian knapweed/ etc.), saltgrass and saltgrass-sacaton, and bottomland grassland (possibly dominated by saltgrass, giant sacaton, dropseed, muhly, burrograss, alkali sacaton, galleta, vinemesquite, and/or tobosa). Typically, the vegetation consists of a shrub/grass mixture characterized by fourwing saltbush and greasewood. Tall, mid-grass, and short grasses are present. Blue grama, foxtail, sand dropseed, spike dropseed, giant dropseed, New Mexico feathergrass and tansymustard are common. When the plant community deteriorates, there is an increase in amounts of shrubs and short grasses (USDA NRCS n.d.).

Pre-treatment monitoring was conducted at these sites on January 30, 2012 as part of a restoration project of non-native phreatophytes scheduled for 2011-2012. Post-treatment monitoring was conducted October 7, 2016. Plot T_1 is located in the northwest corner of the boundary area, plot T_2 sits on the southwest corner and plot T_3 is located approximately in the center of the treatment area (Fig. 3). The project was sponsored by the Valencia Soil and Water Conservation District. Restoration goals are to restore the area for wildlife use, particularly wild turkey habitat, and to remove non-native woody invasive plants (Miller, undated).




Figure 5. Tome project boundary

Tome Site Summary

2016 Tome Site observations: The project area remained the same as observed in 2012. With the exception of the appearance of new salt cedar sprouts and Russian thistle. Plants such as, Vine mesquite, drop seed, Canada wild rye, Kochia, fetid marigold, Mares tail, Aster and Broom snakeweed were observed only in 2016. The area near the river still consisted of a light, multi-tiered understory and a mostly cottonwood over story, as seen in 2012. Much of the area consists of grassy openings. Large downed woody debris and masticated material was present throughout the site. Since monitoring was done so late in the fall, sparse forb and grasses cover may be attributed to seasonal dormancy. The plots were assessed to fall in Hink & Ohmart Structure Classes 2, 4, 5, and 6. Identification of forb, grasses and some shrub species was also impacted by the limited plant identification skills of the monitoring team and by the season

2012 Tome Site observations: The project area is lightly wooded, with most of the wooded area near the river and consisting of a light, multi-tiered understory and a mostly cottonwood over story. It had been treated in the mid-2000s. Much of the area consists of grassy openings. Large downed woody debris and masticated material was present throughout the site. Since monitoring was done so late in the fall, sparse forb and grasses cover may be attributed to seasonal dormancy. The plots were assessed to fall in Hink & Ohmart Structure Classes 2, 4, 5, and 6. Identification of forb, grasses and some shrub species was also impacted by the limited plant identification skills of the monitoring team and by the season

2022 Tome Site Observation: The site was majority open with many large cottonwoods. Some plots were overcrowded with kochia and Russian Olive.

Tome 2012 & 2022 - Observed plant species summary

11.09 Tome						
Vegetation Type/Year	2011		2016		2022	
Graminoids	<i>Distichlis spicata</i>	Inland Saltgrass	<i>Distichlis spicata</i>	Inland Saltgrass	<i>Distichlis spicata</i>	Inland Saltgrass
	<i>Elymus canadensis</i>	Canada Wild Rye	<i>Sporobolus airoides</i>	Alkali Sacaton	<i>Sporobolus airoides</i>	Alkali Sacaton
	<i>Panicum obtusum</i>	Vinemesquite grass	<i>Sporobolus cryptandrus</i>	Sand Dropseed	<i>Sporobolus cryptandrus</i>	Sand Dropseed
	X	Unknown Bunchgrass	X	Unknown Bunchgrass		
Forbs	<i>Machaeranthera canescens</i>	Tansyaster	<i>Bassia prostrata</i>	Kochia	<i>Bassia prostrata</i>	Kochia
			<i>Salsola tragus</i>	Russian Thistle	<i>Salsola spp</i>	Russian Thistle
			<i>Conyza canadensis</i>	Marestail		
			<i>Dyssodia papposa</i>	Fetid Marigold		
Cactus	<i>Opuntia spp</i>	Prickly Pear				
Shrubs	<i>Salix exigua</i>	Coyote Willow	<i>Salix exigua</i>	Coyote Willow	<i>Salix exigua</i>	Coyote Willow
	<i>Artemisia fridiga</i>	Fringed Sagewort	<i>Artemisia fridiga</i>	Fringed Sagewort		
			<i>Gutierrezia sarothrae</i>	Broom Snakeweed		
Trees	<i>Elaeagnus angustifolia</i>	Russian Olive	<i>Elaeagnus angustifolia</i>	Russian Olive	<i>Elaeagnus angustifolia</i>	Russian Olive
	<i>Populus deltoides</i>	Rio Grande Cottonwood	<i>Populus deltoides</i>	Rio Grande Cottonwood	<i>Populus deltoides</i>	Rio Grande Cottonwood
			<i>Tamarix ramosissima</i>	Tamarisk	<i>Tamarix ramosissima</i>	Tamarisk

Figure 6. Species list for the entire project.

New plants observed in 2016 included; Vine mesquite, drop seed, Canada wild rye, Kochia, fetid marigold, Mares tail, Aster and Broom snakeweed. The target species found in 2012, Russian olive, was still present in 2016, as re-sprouts. Salt cedar and Russian thistle, which were not recorded in 2012, appeared in 2016. Identification of forb, grasses and some shrub species was impacted by both the plant identification skills of the monitoring team and by the season.



Figure 7. Tome plots.

Tree Component

The tree component consists of data collected on the 1/10 acre plot Measurements of tree's diameter at breast height (DBH), height, live crown base height, condition (live, sick or dead), and any significant mistletoe damage. We analyze tree density using Trees Per Acre (TPA) and basal density Basal Area Per Acre (BA/AC). All individual trees recorded for this project were *P. deltoides wislizeni*. Wooded species were removed during treatment and this could explain the high density of seedlings, now invading a cleared vegetative strata.




Figure 8. Displays average Trees, Seedlings and Saplings per acre for the entire project




Figure 9. Displays *E. angustifolia* and *S. exigua* seedlings per acre

11.09 Tome			November 2022	
Individual Plot Summary Table				
Macro Plot Name	Total number of sample trees on plot	Growing Stock		
		Number of growing stock sample trees on plot	Trees per Acre	Basal Area per Acre
11.09_1	0	0	0	0.00
11.09_2	8	6	60	90.61
11.09_3	5	4	40	40.31
Total	Total number of sample trees on plot	Number of growing stock sample trees on plot	Average for all Plots	
			TPA	BA/AC
		13.00	10.00	33.33

Table 1. Displays the Stand Table summary for the entire project. Stand tables summarize the tree data collected in all the plots in a readable format for foresters.

Understory Component

As described above, percent ground cover was estimated at each plot within the 1/100th acre subplot. Figures 6 and 7 display the project average cover for each metric. Total aerial cover may exceed 100% due to vegetation stacking on top of each other. Tree canopy remained steady between 2012 and 2016

but increased in 2022. Of interest is the loss of graminoid aerial cover and not seeing a large increase in forb cover, plants that might occupy the same ecological strata as graminoids. There were no major changes in ground cover.




Figure 10. Displays average aerial cover for the entire project




Figure 10. Displays average ground cover for the entire project

Project: Valencia SWCD**Project Unit: Tome 11.09****Plot: 11.09_1**

11.09_1 Aerial & Ground Cover

	Aerial cover (%) of the 1/100-acre plots						
Year	Tree Canopy	Seedlings <5	Saplings 5-15'	Shrubs <5	Shrubs-Saplings 5-15'	Graminoid	Forb
2012	3%	0%	5%	0%	5%	20%	20%
2016	11%	0%	0%	5%	5%	25%	30%
2022	11%	0%	0%	3%	2%	3%	40%

	Ground cover (%) of the 1/100-acre plots					
Year	Litter	Bare soil	Rock	Gravel	Water or wet soil	Plant basal area
2012	59%	1%	0%	0%	0%	n/a
2016	75%	3%	0%	0%	0%	22%
2022	60%	20%	0%	0%	0%	16%

2012 Hink & Ohmart Type: 2 or 6
2016 Hink & Ohmart Type: 4**2016 Modified Hink & Ohmart Type:** 5**2022 Hink & Ohmart Type:** 5**2022 Modified Hink & Ohmart Type:** 5

2012 Comments: masticated/mulched material and CWD present
2016 Comments: tons of kochia, jetty jacks, road crosses N transect line**2022 Comments:** Open area covered in kochia and grasses, with cottonwoods and russian olives scattered sparsely on the edges of the plot.

Project: Valencia SWCD**Project Unit: Tome 11.09****Plot: 11.09_2****11.09_2 Aerial & Ground Cover**

Aerial cover (%) of the 1/100-acre plot							
Year	Tree Canopy	Seedlings <5	Saplings 5-15'	Shrubs <5	Shrubs-Saplings 5-15'	Graminoid	Forb
2012	78%	0%	15%	0%	0%	15%	0%
2016	64%	5%	30%	0%	0%	30%	15%
2022	85%	75%	0%	0%	0%	0%	0%

Ground cover (%) of the 1/100-acre plot						
Year	Litter	Bare soil	Rock	Gravel	Water or wet soil	Plant basal area
2012	73%	12%	0%	0%	0%	n/a
2016	65%	0%	0%	0%	0%	35%
2022	74%	1%	0%	0%	0%	25%

2012 Hink & Ohmart Type: 4
2016 Hink & Ohmart Type: 1 or 3**2016 Modified Hink & Ohmart Type: 1****2022 Hink & Ohmart Type: 4****2022 Modified Hink & Ohmart Type: 2**

2012 Comments: masticated/mulched material and CWD present.**2016 Comments:** lots of Russian olive & salt cedar resprouts, cottonwood seedlings present.**2022 Comments:** Thicket of russian olive understory surrounding plot center. Cottonwoods make up the overstory, lots of litter and slash as well as scattered grasses.

Project: Valencia SWCD**Project Unit: Tome 11.09****Plot: 11.09_3**

11.09_3 Aerial & Ground Cover

Aerial cover (%) of the 1/100-acre plots							
Year	Tree Canopy	Seedlings <5	Saplings 5-15'	Shrubs <5	Shrubs-Saplings 5-15'	Graminoid	Forb
2012	2%	0%	0%	0%	0%	16%	14%
2016	8%	0%	0%	0%	0%	40%	10%
2022	24%	0%	0%	0%	0%	10%	5%

Ground cover (%) of the 1/100-acre plots						
Year	Litter	Bare soil	Rock	Gravel	Water or wet soil	Plant basal area
2012	20%	50%	0%	0%	0%	n/a
2016	35%	40%	0%	0%	0%	25%
2022	30%	60%	0%	0%	0%	10%

2012 Hink & Ohmart Type: 4
2016 Hink & Ohmart Type: 6**2016 Modified Hink & Ohmart Type: 6H****2022 Hink & Ohmart Type: 6****2022 Modified Hink & Ohmart Type: 6H**

2012 Comments: some masticated/mulched material present**2016 Comments:** low diversity of species, bare ground, and old tire tracks over PC.**2022 Comments:** Large open area covered in grasses and litter. Woody debris (logs) scattered around, as well as scattered cottonwoods on the edges.

Next steps (monitoring)

Continuing forward, the goal of the GRGWA/ NMFWR is that all sites will be revisited for post-treatment monitoring in 5-year intervals. It is our intention and expectation that the data collected in these intervals will reflect any significant changes in disturbance and vegetative communities.

References

- Audubon New Mexico. (2013). *Water Matters: Water for New Mexico Rivers*. Albuquerque, New Mexico: Utton Transboundary Resources Center.
- Committee on Riparian Zone Functioning and Strategies for Management, et al. (2002). *Riparian Areas: Functions and Strategies for Management*. Washington, D.C.: National Academy Press.
- Lightfoot, David & Stropki, C. (2012). *Field Manual for Greater Rio Grande Watershed Alliance Riparian Restoration Effectiveness Monitoring*. Albuquerque, NM: SWCA Environmental Consultants.
- New Mexico Department of Game and Fish Conservation Services Division. (2012). *Bridge and Road Construction/Reconstruction Guidelines for Wetland and Riparian Areas*.
- U.S. Climate Data. (2017). *Climate New Mexico*. Retrieved from U.S. Climate Data: <http://www.usclimatedata.com/climate/new-mexico/united-states/3201>
- USDA NRCS. (2016, 8 10). *Web soil Survey*. Retrieved from <https://websoilsurvey.sc.egov.usda.gov/App/HomePage.htm>
- USDA NRCS n.d. (n.d.). *Ecological Site Description Salty Bottomland R042XA055NM*.
- USDA USFS. (1996, September). *Ecology, Diversity, and Sustainability of the Middle Rio Grande Basin*, RM-GTR-268. (D. M. Finch, & J. A. Tainter, Eds.) Fort Collins, Colorado.

Appendix I – Plot Coordinates Table

Plot_Name	Latitude	Longitude
11.09_1	34.7427	-106.7430
11.09_2	34.7412	-106.7430
11.09_3	34.7421	-106.7420

Appendix II - Modified Hink and Ohmart categories, from NMRAM

The following is pages 39-41 in Muldavin et al.'s 2014 NMRAM for Montane Riverine Wetlands v 2.0 Manual (draft, not yet published)

Vegetation Vertical Structure Type Definitions for NMRAM

Multiple-Story Communities (Woodlands/Forests)



Type 1 – High Structure Forest with a well-developed understory.

Tall mature to intermediate-aged trees (>5 m [>15 feet]) with canopy covering >25% of the area of the community (polygon) and understory layer (0-5 m [0-15 feet]) covering >25% of the area of the community (polygon). Substantial foliage is in all height layers. (This type incorporates Hink and Ohmart structure types 1 and 3.) Photograph on Gila River by Y. Chauvin, 2012.



Type 2 –Low Structure Forest with little or no understory.

Tall mature to intermediate-aged trees (>5 m [>15 feet]) with canopy covering >25% of the area of the community (polygon) and understory layer (1-5 m [3-15 feet]) covering <25% of the area of the community (polygon). Majority of foliage is over 5 m (15 feet) above the ground. (This type incorporates Hink and Ohmart structure types 2 and 4.) Photograph on Diamond Creek by Y. Chauvin, 2012.

Single-story Communities (Shrublands, Herbaceous and Bare Ground)



Type 5 –Tall Shrub Stands.

Young tree and shrub layer only (15-5 m [4.5-15 feet]) covering >25% of the area of the community (polygon). Stands dominated by tall shrubs and young trees, may include herbaceous vegetation underneath the woody vegetation. Photograph on San Francisco River by Y. Chauvin, 2012.




Type 6S-Short Shrub Stands.

Short stature shrubs or very young shrubs and trees (up to 1.5 m [up to 4.5 feet]) covering >10% of the area of the community (polygon). Stands dominated by short woody vegetation, may include herbaceous vegetation underneath the woody vegetation. Photograph on Lower Pecos River by E. Lindahl, 2008.



Type 6W-Herbaceous Wetland.

Herbaceous wetland vegetation covering >10% of the area of the community (polygon). Stands dominated by obligate wetland herbaceous species. Woody species absent, or <10% cover. Photograph of *Carex nebrascensis* meadow on upper Rio Santa Barbara by Y. Chauvin, 2009.

**Type 6H- Herbaceous.**

Herbaceous vegetation covering >10% of the area of the community (polygon). Stands dominated by herbaceous vegetation of any type except obligate wetland species. Woody species absent or <10% cover. Photograph on Diamond Creek by Y. Chauvin, 2012.

**Type 7-Sparse Vegetation/Bare Ground.**

Bare ground, may include sparse woody or herbaceous vegetation, but total vegetation cover <10%. May be natural in origin (cobble bars) or anthropogenic in origin (graded or plowed earth) Photograph on Lower Gila River by Y. Chauvin, 2012.

Appendix III – Sample Datasheet

2011 Datasheet with original Hink & Ohmart

GRGWA 2011 Revisit Data Sheet

Project: _____
 Project Unit: _____
 Plot Number: _____
 Lat (dd.dddd): _____ Long (ddd.dddd): _____ Elevation: _____ ft

Date: _____
 Time: _____

Plot size: 1/100th ac for understory ("small plot")
 1/10th ac for overstory

Aerial cover										Cover % - Taken from/within small plot only					
Tree canopy (use densiometer facing out at 11'9" flags)	Seedlings <5' (estimate aerial cover)	Saplings 5-15' (estimate aerial cover)	Shrubs <5' (estimate aerial cover)	Shrubs 5-15' (estimate aerial cover)	Graminoid (estimate aerial cover)	Forb (estimate aerial cover)	Litter (estimate ground cover)	Bare soil (estimate ground cover)	Rock (estimate ground cover)	Gravel (estimate ground cover)	Water or wet soil (estimate ground cover)	Plant Basal Area (est. ground cover)			

Hink & Ohmart structural class for entire 1/10th ac plot (unmodified, see back):

Hink & Ohmart modified structural class for entire 1/10th acre plot (see NMRM)

Species Observed in 1/10th ac plot (scientific name, common name, or USDA PLANTS code)

Grasses	Forbs	Shrubs	Trees

Photopoints needed (with whiteboard):


- PC showing whiteboard with name clearly legible
- North facing Center – 66'
- PC north to 11'9"
- PC east to 11'9"
- PC south to 11'9"
- PC west to 11'9"

Comments/Observations:

Unmodified Hink & Ohmart (courtesy of SWCA)


2022 Sample datasheet

GRGWA Plot Description (1 of 2)

Observer: _____ Recorder: _____ Latitude (dd.dddddd): _____ Longitude (ddd.ddddd): _____ Elevation (ft): _____			Administrative Unit: _____ Project Unit: _____ Macroplot: _____ Date (DD/MM/YYYY): _____ Time: _____																																																																																																																														
Macroplot Sizes <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Size (Acres)</td> <td style="width: 20%;">1/100</td> <td style="width: 20%;">1/10</td> </tr> <tr> <td>Radius (Feet, Decimal Feet)</td> <td>11.78</td> <td>37.24</td> </tr> <tr> <td>Radius (Feet, Inches)</td> <td>11' 9"</td> <td>37' 3"</td> </tr> </table>			Size (Acres)	1/100	1/10	Radius (Feet, Decimal Feet)	11.78	37.24	Radius (Feet, Inches)	11' 9"	37' 3"	Hill Slope (where steepest): _____ % Aspect (circle one): N E S W Aspect azimuth: _____ ° Mag Declination: _____ °																																																																																																																					
Size (Acres)	1/100	1/10																																																																																																																															
Radius (Feet, Decimal Feet)	11.78	37.24																																																																																																																															
Radius (Feet, Inches)	11' 9"	37' 3"																																																																																																																															
Photo Azimuths: (1) of whiteboard at PC. (1) from 75 feet N looking south to PC (4) from PC in all four cardinal directions; (1) from each Brown's transect looking toward PC. ORDER TAKEN: _____			<div style="text-align: center;">  </div> Describe Witness Tree(s): USE NATIVE TREES ONLY **Draw location of tree on plot** Color of Flagging Used: _____																																																																																																																														
Tree Canopy Cover (%) (densiometer) <div style="border: 1px solid black; height: 40px; margin-top: 5px;"></div>			Comments/Description of Plot: 																																																																																																																														
Hink & Ohmart Dominant Structural Class Original: Modified:																																																																																																																																	
Soil Texture (4 locations) North: _____ East: _____ South: _____ West: _____			**SMALL PLOT INCLUDES ALL SEEDLINGS OR SAPLINGS <5 INCHES DBH/DRC.** <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th rowspan="3">Species</th> <th rowspan="3">Condition (Live, Dead, Sick)</th> <th colspan="5">Small Plot (1/100th Acre only) - Tree Regen, Shrubs & Cacti</th> <th rowspan="3">Species</th> <th rowspan="3">Condition (Live, Dead, Sick)</th> <th colspan="5">Small Plot (1/100th Acre only) - Tree Regen, Shrubs & Cacti</th> </tr> <tr> <th colspan="5">Height classes—Seedlings (feet)</th> <th colspan="5">Diameter classes—Saplings (inches)</th> </tr> <tr> <th>> 0 - 0.5'</th> <th>> 0.5 - 1.5'</th> <th>> 1.5' - 2.5'</th> <th>> 2.5' - 3.5'</th> <th>> 3.5' - 4.5'</th> <th>> 0 - 1"</th> <th>> 1-2"</th> <th>> 2-3"</th> <th>> 3-4"</th> <th>> 4-5"</th> </tr> </thead> <tbody> <tr><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td></tr> <tr><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td></tr> <tr><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td></tr> <tr><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td></tr> <tr><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td></tr> <tr><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td><td> </td></tr> </tbody> </table>			Species	Condition (Live, Dead, Sick)	Small Plot (1/100th Acre only) - Tree Regen, Shrubs & Cacti					Species	Condition (Live, Dead, Sick)	Small Plot (1/100th Acre only) - Tree Regen, Shrubs & Cacti					Height classes—Seedlings (feet)					Diameter classes—Saplings (inches)					> 0 - 0.5'	> 0.5 - 1.5'	> 1.5' - 2.5'	> 2.5' - 3.5'	> 3.5' - 4.5'	> 0 - 1"	> 1-2"	> 2-3"	> 3-4"	> 4-5"																																																																																										
Species	Condition (Live, Dead, Sick)	Small Plot (1/100th Acre only) - Tree Regen, Shrubs & Cacti						Species	Condition (Live, Dead, Sick)	Small Plot (1/100th Acre only) - Tree Regen, Shrubs & Cacti																																																																																																																							
		Height classes—Seedlings (feet)								Diameter classes—Saplings (inches)																																																																																																																							
		> 0 - 0.5'	> 0.5 - 1.5'	> 1.5' - 2.5'	> 2.5' - 3.5'	> 3.5' - 4.5'	> 0 - 1"			> 1-2"	> 2-3"	> 3-4"	> 4-5"																																																																																																																				

New Mexico Forest and Watershed Restoration Institute

Plot Description Version: 4/3/2018, km



Precisions:
 Slope: ±5 percent
 Vegetation cover ±1 class estimation or ±10%

GRGWA Plot Description (2 of 2)

[illegible]

GROUND COVER (%) (ENTIRE 1/10th acre plot) (must total 100 %)							
Plant basal	Bole	Litter	Bare soil	Rock (>2.5in)	Gravel (< 2.5 in)	Water, Wet Soil	Total (%)

Comments on Species Composition and/or Ground Cover:

Appendix IV – Fuels Transect Data Sheet

GRGWA Surface Fuels

Sheet 1 of 1: Fine Woody Debris—Coarse Woody Debris

Observer _____ Recorder _____	Administrative Unit: _____ Project Unit: _____ Macroplot: _____ Date (DD/MM/YYYY): _____ Time: _____
--	---

1-hour Transect Length - 6'	10-hour Transect Length - 6'
100-hour Transect Length - 35'	1000-hour Transect Length - 60'

Class	Diameter (in)	
FWD	1-hr	0 to 0.25
	10-hr	0.25 to 3.0
	100-hr	3.0 to 3.0
CWD	1000-hr and greater	3.0 and greater



The diagram shows a horizontal transect line with two sampling cylinders. The left cylinder is labeled 'Duff layer sampling cylinder' and the right cylinder is labeled 'Vegetation sampling cylinder'. The distance between the cylinders is 25.0. The distance from the left cylinder to the right edge of the transect is 25.0. The distance from the right cylinder to the right edge of the transect is 25.0. The total length of the transect is 75.0. The diagram also shows a scale bar from 0 to 100 feet.

Fine Woody Debris (1, 10, 100 hr fuels)	Transect	Azimuth	Slope	1 - Hr Count	10 - Hr Count	100 - Hr Count	Comment
1, 10, 100 hr fuels	1						
	2						

Coarse Woody Debris (1000 hr fuels)	Transect	Slope	Log No.	Log Diameter	Decay Class	Comment
Coarse Woody Debris (1000 hr fuels)						

Litter & Duff	Transect 1	45'	75'	Transect 2	45'	75'
Litter & Duff	Litter Depth (in)			Litter Depth (in)		
	Duff Depth (in)			Duff Depth (in)		
	Comments?			Comments?		

Precisions: Diameter: ± 0.5 in ; decay class ± 1 class ; Slope ± 5 percent

Decay Class Description

- 1 All bark is intact. All but the smallest twigs are present. Old needles probably still present. Hard when kicked
- 2 Some bark is missing, as are many of the smaller branches. No old needles still on branches. Hard when kicked
- 3 Most of the bark is missing and most of the branches less than 1 in. in diameter also missing. Still hard when kicked
- 4 Looks like a class 3 log but the sapwood is rotten. Sounds hollow when kicked and you can probably remove wood from the outside with your boot. Pronounced sagging if suspended for even moderate distances
5. Entire log is in contact with the ground. Easy to kick apart but most of the piece is above the general level of the adjacent ground. If the central axis of the piece lies in or below the duff layer then it should not be included in the CWD sampling as these pieces act more like duff than wood when burned.

Appendix V – Photo Pages

10-Year Photo Comparison for 11.09 Tome, 3 plots

2011/2012 photos: taken January 31, 2012 by Joe Zebrowski

2016/2017 photos: taken November 6, 2016 by Kathryn Mahan

2022 photos: taken November by Alex Makowicki / Clay Goetsch

Contact:

Alex Makowicki, Ecological Monitoring Technician

alexmakowicki@nmhu.edu



11.09_1C Facing center from north at 66' (2012)



11.09_1C facing center from north at 66' (2016)



11.09_1C facing center from north at 66' (2022)



11.09_1N Facing north from center at 11.8' (2012)



(2016)

11.09_1N facing north from center at 11.8'



(2022)

11.09_1N facing north from center at 11.8'



11.09_1E facing east from center at 11.8' (2012)



11.09_1E facing east from center at 11.8' (2016)



11.09_1E facing east from center at 11.8' (2022)



11.09_1S facing south from center at 11.8'



11.09_1S facing south from center at 11.8'

(2016)



11.09_1S facing south from center at 11.8'

(2022)



11.09_1W facing west from center at 11.8' (2012)



(2016)

11.09_1W facing west from center at 11.8'



(2022)

11.09_1W facing west from center at 11.8'



11.09_2C facing center from north at 66' (2012)




11.09_2C facing center from north at 66' (2016)



11.09_2C facing center from north at 66' (2022)



11.09_2N facing north from center at 11.8' (2012)



(2016)

11.09_2N facing north from center at 11.8'



(2022)

11.09_2N facing north from center at 11.8'



11.09_2E facing east from center at 11.8' (2012)



11.09_2E facing east from center at 11.8' (2016)



11.09_2E facing east from center at 11.8' (2022)



11.09_2S facing south from center at 11.8' (2012)



(2016)

11.09_2S facing south from center at 11.8'




(2022)

11.09_2S facing south from center at 11.8'



11.09_2W facing west from center at 11.8' (2012)



(2016)

11.09_2W facing west from center at 11.8'



(2022)

11.09_2W facing west from center at 11.8'



11.09_3C facing center from north at 66' (2012)



11.09_3C facing center from north at 66' (2016)



11.09_3C facing center from north at 66' (2022)



11.09_3N facing north from center at 11.8' (2012)




(2016)

11.09_3N facing north from center at 11.8'



(2022)

11.09_3N facing north from center at 11.8'



11.09_3E facing east from center 11.8' (2012)



11.09_3E facing east from center at 11.8' (2016)



11.09_3E facing east from center at 11.8' (2022)



11.09_3S facing from center at 11.8' (2012)



(2016)

11.09_3S facing south from center at 11.8'



11.09_3S facing south from center at 11.8' (2022)



11.09_3W facing west from center at 11.8' (2012)



11.09_3W facing west from center at 11.8' (2016)



11.09_3W facing west from center at 11.8' (2022)