

Grants SE Project 16.12


Post-treatment Monitoring Report

2021



Prepared by

Kathryn R Mahan, Ecological Monitoring Specialist & Alex Makowicki, Ecological Monitoring Technician,
New Mexico Forest and Watershed Restoration Institute
for the Greater Rio Grande Watershed Alliance



Lava SWCD

Contents

Acronyms and Abbreviations	3
Purpose of Report	4
Ecological Context of Bosque Restoration	4
Monitoring and Field Methods	5
Low intensity Field Methods	5
High-intensity Field Methods	6
Personnel Involved	6
Grants SE Project	7
Discussion	12
References	13
Appendix I - Photopoint Table	15
Appendix II - Photos	18
Appendix III – Current monitoring methods available	26
Appendix IV - Modified Hink and Ohmart categories, from NMRAM	27

Acronyms and Abbreviations

Acronym, Abbreviation, or Term	Explanation or Definition as used by NMFWR
AGL	above ground level; GIS term
BBIRD plots	Breeding Biology Research and Monitoring Database, larger circular plot types
BEMP plots	Bosque Ecosystem Monitoring Program, small rectangular plot types
FEAT	Fire Ecology Assessment Tool
FFI	FEAT/ FIREMON Integrated
FIREMON	Fire Effects Monitoring and Inventory System
FSA	Farm Service Agency, a department of the USDA
GIS	Geographic Information Systems
GRGWA	Greater Rio Grande Watershed Alliance
LIDAR	Light detecting and ranging, a remote sensing technique using light to gather elevation data
NAIP	National Agriculture Imagery Program (aerial imagery)
NDVI	Normalized Difference Vegetation Index; GIS term for a band ratio of the visible red and the near infrared spectral bands and is calculated using the following formula: $(NIR - Red)/(NIR + Red)$
NHNM	Natural Heritage New Mexico
NMDGF	New Mexico Department of Game and Fish
NMED SWQB	New Mexico Environment Department Surface Water Quality Bureau
NMFWR	New Mexico Forest and Watershed Restoration Institute
NMHU	New Mexico Highlands University
NMRAM	New Mexico Rapid Assessment Method, version 2.0
NRCS	Natural Resource Conservation Service
PC	Plot center
RGIS	Resource Geographic Information System
SWCD	Soil and Water Conservation District
TIFF	Tagged image file format
USDA	United States Department of Agriculture
USGS	United States Geological Survey
WQCC	Water Quality Control Commission
WSS	Web Soil Survey, a soils database of the NRCS

Purpose of Report

This report covers the low-intensity pre-treatment vegetation monitoring assessment performed on a non-native phreatophyte removal project submitted for a drainage basin adjacent to the Rio San Jose channel in the city of Grants, NM, to the Greater Rio Grande Watershed Alliance in 2016. Following a discussion of the ecological context, and our monitoring methods, we present pertinent background, observations, and assessment results for the project.

Ecological Context of Bosque Restoration

Neither the challenges nor the importance of working in the bosque and other riparian areas in New Mexico today should be underestimated. According to the New Mexico Department of Game and Fish Conservation Division, wetlands and riparian areas comprise approximately 0.6 percent of all land in New Mexico (2012). Despite this small percentage, estimates of New Mexican vertebrate species depending on wetland and riparian habitat for their survival ranges from 55% (New Mexico Department of Game and Fish Conservation Services Division, 2012) to 80% (Audubon New Mexico, 2013). These areas also provide flood mitigation, filtration of sediment and pollutants, and water for a variety of purposes including groundwater recharge (Audubon New Mexico, 2013). In addition, native vegetation such as cottonwoods have cultural significance to many communities.

As much as these areas are disproportionately important to ecosystems and human communities, they are equally disproportionately impacted by disturbance. Anthropogenic impacts with major consequences for our riparian areas include dams, reservoirs, levees, channelization, acequias and ditches, jetty jacks, riprap and Gabion baskets, urbanization, removal of native phreatophytes, grazing by domestic livestock, excessive grazing pressure by native ungulate populations absent natural predation cycles, beaver removal, logging, mining, recreation, transportation, introduction and spread of invasive exotic species, groundwater extraction, altered fire and flood regimes drought and climate change (Committee on Riparian Zone Functioning and Strategies for Management, et al., 2002). Statewide, it is estimated that as much as 90% of New Mexico's historical riparian areas have been lost (Audubon New Mexico, 2013), and approximately 39% of our remaining perennial stream miles are impaired (New Mexico Department of Game and Fish Conservation Services Division, 2012).

New Mexico is fortunate enough to have the Middle Rio Grande Bosque, the largest remaining bosque in the Southwest (USDA USFS, 1996). However, over the past two decades, the number of fires in the bosque has been increasing. Historically, the primary disturbance regime in the bosque has been flooding, not fire, which means the system is not fire-adapted. In fact, native species like cottonwood resprout from their roots after floods and need wet soils to germinate from seed. Flooding also promotes decomposition of organic material and keeps the soil moist which reduces the likelihood of fire. Today, overbank flow is uncommon in many areas of the Rio Grande due to the heavy alteration of the channel and flow regimes (two obvious examples are the structures defining the upper and lower extent of the Middle Rio Grande: Cochiti Dam and Elephant Butte Reservoir). This has led to low fuel moisture content and high fuel loads, as well as increased human presence in the riparian area. As a result, bosque fires are more common and more severe: they kill cottonwoods and other native species, creating spaces which are filled by non-native species such as salt cedar, Russian olive, Siberian elm, and Tree-of-Heaven. We are constantly learning more about how these species can exploit and encourage a riparian fire regime, in addition to many other changes they bring to ecosystems.

Efforts geared toward the removal of these nonnative species can help to reduce fire risk, preserve native vegetation, and be part of a larger effort to restore the bosque and the watershed as a whole to a more natural and functional ecosystem. The Greater Rio Grande Watershed Alliance (GRGWA) has been working on these issues with a variety of collaborating organizations and agencies within the Rio Grande basin for several years. Since 2013, the New Mexico Forest and Watershed Restoration Institute (NMFWRRI) has been working with GRGWA and the Claunch-Pinto Soil and Water Conservation District (SWCD) to begin construction of a geodatabase for all of GRGWA's non-native phreatophyte removal projects as well as to perform the formal pre- and post-treatment monitoring, utilizing the field methods explained below as well as LIDAR analysis where appropriate and available.

Monitoring and Field Methods

Low intensity Field Methods

Low intensity pre-treatment vegetation monitoring was done using an adapted version of the biotic portion of the New Mexico Rapid Assessment Method (NMRAM), v 2.1, updating recommendations made in the Field Manual for Greater Rio Grande Watershed Alliance (GRGWA) Riparian Restoration Effectiveness Monitoring and the GRGWA Monitoring Plan, developed by Lightfoot & Stropki of SWCA Environmental Consultants in 2012. (For a brief overview of both low and high intensity monitoring methods used by the NMFWRRI on GRGWA projects, please see Appendix III.)

For those not familiar, NMRAM was developed by the New Mexico Environment Department Surface Water Quality Bureau Wetlands Program and Natural Heritage New Mexico as a "cost effective, yet consistent and meaningful tool" (Muldavin, 2011) for wetland ecological condition assessment in terms of anthropogenic disturbance as negatively correlated with quality and functionality. The portions of NMRAM we utilized are Level 2 "semi-quantitative" field measurements taken at less detail than plot level (Muldavin, 2011).

Measurements taken included relative native plant community composition, vegetation horizontal patch structure, vegetation vertical structure, native riparian tree regeneration, and invasive exotic plant species cover. The underlying method for these biotic assessments was a version of the 1984 Hink and Ohmart vertical structure classification system, modified for use in the NMRAM for Montane Riverine Wetlands version 2.0 (see Appendix IV). First, vegetation communities were mapped out by patch (polyon) according to the Hink and Ohmart system. Next, the presence of (state-listed) invasives, wetland species, and the two dominant species in each strata ("tree" >15 ft, "shrub" 4.5-15 ft, and "herbaceous" <4.5 ft) were recorded for each plant community. The native/exotic ratio in each of the patches was scored and weighted based on the percent of the project area each patch comprised. These scores were then combined with the additional biotic metrics of vertical and horizontal diversity, native tree regeneration, and overall (listed) invasive presence. The NMRAM rating system is based, on all levels, on a scale of 1 to 4, where 4 is considered excellent condition, 3 good, 2 fair, and 1 poor.


We also assessed soil surface condition, which is a metric typically included in the abiotic section of the NMRAM, as well as the presence of surface fuels, which is not part of the NMRAM. Unlike the other 6 metrics we used, surface fuels were recorded on a rating scale from 0 to 1.0 where 1.0 is a continuous fuel matrix.

Photopoints were established to capture images where vegetation shifts were observed and/or at representative locations throughout the site. Waypoints were marked with a Garmin GPS unit and named sequentially by site. Photos were taken facing north, east, south and west at each point.

Prior to entering the field, we created a map with the project boundaries as provided by GRGWA. We combined these polygons with recent aerial imagery and identified relevant roads and other landscape features. Once on the ground, the vegetation community polygons (as determined by the modified Hink and Ohmart classification system) were hand-drawn onto this map and served as the basis for other biotic metric assessments. Upon return to the office, this polygon map and the photopoints were digitized by the monitoring technician and/or specialist.

High-intensity Field Methods

High-intensity monitoring was also done, in part, on this site. We used an adapted Bosque Ecosystem Monitoring Program (BEMP) style plot. These are 16 x 98-foot rectangles, placed approximately parallel to the river. Within these plots, we measure canopy and species, and vegetation and ground cover. We also used Brown's transects to measure surface fuels.



Personnel Involved

2021 New Mexico Forest and Watershed Restoration Institute Monitoring Team:

- Kathryn R Mahan, Monitoring Program Manager
- Carmen Briones, Crew Logistics Support/ Assistant Manager
- Raymundo Melendez, Monitoring Technician
- Alex Makowicki, Monitoring Technician

2021 New Mexico Forest and Watershed Restoration Institute GIS Team:

- Patti Dappen, GIS Program Manager
- Katie Withnall, GIS Specialist

Other persons contacted:

- Fred Rossbach, Field Coordinator, Greater Rio Grande Watershed Alliance
- Larry Winn, Lava/McKinley Soil and Water Conservation District

Grants SE Project

Project 16.12 is located in a meadow area/drainage basin adjacent to the Rio San Jose channel and drainage. The area is on private and municipal owned lands on the southeast side of the city of Grants, NM.

The average annual precipitation in Grants, NM is 10.5 inches. The average high temperature is 91° F in July, and the average low is 15° F in December and January. (U.S. Climate Data, 2017) According to the NRCS Web Soil Survey, the project area is comprised of 71% Warm Springs loam, 0 to 2 percent slopes, 22% Venadito clay loam, 0 to 1 percent slopes, and 7.2% Viuda-Penistaja-Rock outcrop complex, 1 to 10 percent slopes. Ecological sites within this project include R035XA112NM Loamy, R035XA119NM Clayey Bottom, and R036XB007NM Malpais (USDA NRCS, 2016).


The Loamy ecological site typically supports a grassland state dominated by blue grama, western wheatgrass, galleta, ring muhly, dropseeds, and/or threeawns. It can also be found in a piñon-juniper invaded state (dominated by piñon, juniper, and blue grama), a grass/succulent-mix state (dominated by blue grama, cholla and prickly pear), a shrub-dominated state (dominated by rabbitbrush or horsebrush and blue grama), as well as a bare state with sparse grass (USDA NRCS n.d.).

The Clayey Bottomland ecological site typically supports a grassland state dominated by western wheatgrass, blue grama, galleta, and alkali sacaton. It can also be found in a shrub-dominated state where dominant vegetation is rabbitbrush, fourwing saltbush, galleta and other grasses, as well as a bare state with sparse annual vegetation (USDA NRCS n.d.).

The Malpais ecological site typically supports a grassland state dominated by blue grama and sideoats grama. Other common vegetation includes western wheatgrass, little bluestem, spike muhly, black grama, galleta, New Mexico feathergrass, alkali sacaton, winterfat, fourwing saltbush, broom snakeweed, and scattered piñon and juniper. In a deteriorated state, the grass community may become sod-bound, dominated by blue grama sod, threeawns, wolfstail and snakeweed (USDA NRCS n.d.).

Post-treatment monitoring was conducted at this 50-acre site on December 9th, 2021 as part of a restoration project targeting non-native phreatophytes conducted in 2016-2017. The area is on private and municipal lands on the southeast side of Grants, NM. The project is bordered on the west by urban development, on the north by the railroad right-of-way, on the northeast by the Rio San Jose channel, on the east by undeveloped Pueblo of Acoma lands and on the south by the Interstate 40 frontage road. The Lava Soil and Water Conservation District (LSWCD) sponsored the project. The project is an initial treatment to remove and treat non-native phreatophyte species as well as down, dead fuels. Salt cedars, Russian olive, and Siberian elm trees will be treated and removed with extraction and mastication; however, large Siberian elms used as shade trees near urban development will be left. Restoration goals include improving awareness of bosque restoration efforts, restoring the riparian area to a more natural state, promoting native plant species, and reducing fire hazard.

16-12 SE Grants



Project 16-12 SE Grants is located in a meadow area/drainage basin adjacent to the Rio San Jose channel and drainage. The area is on private and municipal owned lands on the southeast side of the city of Grants, NM. The project is an initial treatment to remove and treat non-native phreatophytes such as salt cedar, Russian olive, Siberian elm trees and down, dead fuels.



Figure 1. Project 16.12 in geographic context.

16-12 SE Grants



Figure 2. 16.12 Grants project outline.

Access to the project is possible through various point including a frontage road and hotel parking lots. There are multiple landowners, fence lines, roads, powerlines and structures; construction debris and trash were present throughout the site at the time of the site visits. Exotic species observed during the monitoring visit included salt cedar (of all age classes) and *phragmites* reed grasses. Native plants observed included fourwing saltbush, inland saltgrass, dropseed grass, and bush seepweed (*Suaeda nigra*). Species density varied within the project and was especially sparse around the playas/salt flats.


Table 1. NMRAM Scores for 16.12.

Metrics for 16.12 (December 9, 2021)	2021 Score	2016 Score
Relative Native Plant Community Composition	2	1
Vegetation Horizontal Patch Structure	2	4
Vegetation Vertical Structure	1	2
Native Riparian Tree Regeneration	1	1
Exotic Invasive Plant Species Cover	1	1
Project Biotic Score (based on above ratings)	1.5	1.8
Project Biotic Rating	D/Poor	C/Fair
Soil Surface Condition	2	4
Surface Fuels	0.70	0.2

Low scores came in Vegetation Horizontal Structure and Vegetation Vertical Structure due to a lack of vegetation structure diversity. The Vegetation Vertical Structure score dropped in 2021 due to the dominance of herbaceous vegetation on land that was previously dominated by tall shrubs, it should be noted that the loss of some of the tall shrub vegetation structure can be attributed to the removal of invasive woody plants such as Tamarisk and Russian Elm. Overall, the site received a “D” or “Poor”, a 1.5 out of 4

In 2016 a BEMP plot was taken to include more detailed monitoring. In 2021, time did not allow for the taking of a BEMP plot.

Figure 3. 16.12 Grants project vegetation polygons.



Discussion

We would like to clarify that we are adapting these NMRAM metrics for our own purposes. That is, we are using them both inside and outside their intended site ranges, including on larger sites (NMRAM is designed to handle a site around 100 x 200 meters), sites further from the river (NMRAM is currently in use primarily for assessing riverine wetlands), and sites defined by exotic vegetation presence rather than hydrologic boundaries and upland vegetation indicators/apparent wetland extent. Site delineation and size is likely to be variable for a number of other reasons, including landowner participation, available funds, proposals received from contractors, etc – many of which cannot be directly correlated to site disturbance or ecological function. For this reason, we do not use the entire NMRAM assessment, or place confidence in the weighted score roll-ups that are typically part of an NMRAM report. Should one be interested, rationale for the weighting in the NMRAM score roll-up can be found in the yet-to-be-published field manual for version 2.1. For more information, contact Maryann McGraw of the NMED or NMFWR.

While we provide a biotic site score and rating for your reference, we recommend comparisons be done with individual metrics from pre-treatment and post-treatment assessment from the same site, rather than across multiple sites. Also of note is that statistical analysis is not appropriate for NMRAM, or other low intensity, rapid field methods.

Please note that should the project area change significantly from what was originally proposed and monitored, all metrics will lose some amount of confidence on comparison as it is impractical to re-examine the original site assessment scores using new boundaries. This is an issue of concern of which GRGWA should be aware. We recommend that GRGWA attempt to minimize alterations in project boundaries once pre-treatment monitoring data has been approved for collection. Another, somewhat alternative, recommendation is that the initial monitoring regime include high-intensity modified BEMP-type plots which could be repeated in their exact initial locations, allowing collection of comparable data regardless of boundary change. We recognize that this is not always practical: boundaries change for a number of reasons and time and cost constraints can necessitate the sole use of a rapid assessment method for monitoring. We have reason to hope our outlined assessment method will still be a satisfactory indicator for site function improvement or degradation primarily because metrics in rapid assessment methods such as this are set up to have relatively low sensitivities (i.e. for a change to be reflected in the metrics, either positive or negative, disturbance on site has to be significantly altered).

From here on out, the goal of the GRGWA/ NMFWR is that all sites will be revisited for post-treatment monitoring in 5-year intervals. It is our intention and expectation that the data collected in these intervals will reflect any significant changes in disturbance and ecological function of the site.

References

- Audubon New Mexico. (2013). *Water Matters: Water for New Mexico Rivers*. Albuquerque, New Mexico: Utton Transboundary Resources Center.
- Brown, J. K. (1974). *Handbook for Inventorying Downed Woody Material*, USDA Forest Service General Technical report INT-16. *Handbook for Inventorying Downed Woody Material*. Ogden, Utah: USDA Forest Service Intermountain Forest and Range Experiment Station.
- Bureau of Land Management. (2006). *Grazing Management Processes and Strategies for Riparian-Wetland Areas, TR 1737-20*.
- Claunch-Pinto Soil and Water Conservation District on behalf of the Greater Rio Grande Watershed Alliance. (2015). *Request for Proposals for Greater Rio Grande Watershed Alliance Riparian Restoration Projects*. Mountainair, NM: Claunch-Pinto Soil and Water Conservation District.
- Committee on Riparian Zone Functioning and Strategies for Management, et al. (2002). *Riparian Areas: Functions and Strategies for Management*. Washington, D.C.: National Academy Press.
- Lightfoot, D. &. (2012). *Greater Rio Grande Watershed Alliance Riparian Restoration Effectiveness Monitoring Plan*. Albuquerque, NM: SWCA Environmental Consultants.
- Lightfoot, David & Stropki, C. (2012). *Field Manual for Greater Rio Grande Watershed Alliance Riparian Restoration Effectiveness Monitoring*. Albuquerque, NM: SWCA Environmental Consultants.
- Lizarazo, I., & Elsner, P. (2009). Fuzzy segmentation for object-based image classification. *International Journal of Remote Sensing*, 30.
- Muldavin, E. B. (2011). *New Mexico Rapid Assessment Method: Montaine Riverine Wetlands*. Version 1.1. Final report to the New Mexico Environment Department, Surface Water Quality Bureau. 90 pp. and appendices.
- New Mexico Department of Game and Fish Conservation Services Division. (2012). *Bridge and Road Construction/Reconstruction Guidelines for Wetland and Riparian Areas*.
- U.S. Climate Data. (2017). *Climate New Mexico*. Retrieved from U.S. Climate Data: <http://www.usclimatedata.com/climate/new-mexico/united-states/3201>
- USDA NRCS. (2016, 8 10). *Web soil Survey*. Retrieved from <https://websoilsurvey.sc.egov.usda.gov/App/HomePage.htm>
- USDA NRCS n.d. (n.d.). *Ecological Site Description Clayey Bottomland R035XA119NM*.
- USDA NRCS n.d. (n.d.). *Ecological Site Description Loamy R035XA112NM*.
- USDA NRCS n.d. (n.d.). *Ecological Site Description Malpais R036XB007NM*.
- USDA USFS. (1996, September). *Ecology, Diversity, and Sustainability of the Middle Rio Grande Basin*, RM-GTR-268. (D. M. Finch, & J. A. Tainter, Eds.) Fort Collins, Colorado.

Appendix I - Photopoint Table

Name	Latitude	Longitude
16.12_1_NESW	35.1220	-107.8230
16.12_2_NESW	35.1311	-107.8230
16.12_PC_NESW	35.1295	-107.8240

Photo Name	Date/Time	Photo Number	Latitude	Longitude	Direction	Foreground Description	Background Description	Camera	Crew	Notes
16.12_PC_NESW	12/8/2021 9:44am	100-0381	35.12949 959	-107.82438 04	PC	-	-	Olympus red	AM, CB, RM	On bareground with Motel 6, next to dirt road.
16.12_PC_NESW	12/8/2021 9:44am	100-0383	35.12949 959	-107.82438 04	N	Tumbleweed, bareground @13' @314°	Fourwing salt bush @153' @360°	Olympus red	AM, CB, RM	Bareground, dirt road, salt cedar, fourwing salt cedar, traintrack, mountain range.
16.12_PC_NESW	12/8/2021 9:44am	100-0384	35.12949 959	-107.82438 04	E	Fourwing salt bush @10' @90°	Salt cedar @544' @90°	Olympus red	AM, CB, RM	Dirt road, fourwing salt bush, grassy area, salt cedar, buildings, mountain range.
16.12_PC_NESW	12/8/2021 9:44am	100-0385	35.12949 959	-107.82438 04	S	Fourwing salt bush @20' @180°	Dollar tree store @1036' @180°	Olympus red	AM, CB, RM	Bareground, fourwing salt bush, Dollar store, Walmart, common reed, rushes.

16.12 _PC_N ESW	12/8/2 021 9:44am	100- 0389	35.12949 959	- 107.82438 04	W	Fourwing salt bush @6' @270°	Salt cedar @184' @270°	Olympus red	AM, CB, RM	Mulching, bareground, tumbleweed, common reed, salt cedar, asian buffet.
16.12 _1_NE SW	12/8/2 021 8:59am	100- 0375	35.12199 482	- 107.82351 06	N	Fourwing salt bush @10' @360°	Salt cedar @314' @360°	Olympus red	AM, CB, RM	Tumbleweed, bareground, saltcedar, fence, buildings and mountain range.
16.12 _1_NE SW	12/8/2 021 8:59am	100- 0376	35.12199 482	- 107.82351 06	E	Mulching/tu mbleweed @19' @90°	Salt cedar @180' @90°	Olympus red	AM, CB, RM	Tumbleweed, mulching, salt cedar, mountain range
16.12 _1_NE SW	12/8/2 021 8:59am	100- 0379	35.12199 482	- 107.82351 06	S	Mulching/tu mbleweed @67' @180°	Hwy I-40 @546' @180°	Olympus red	AM, CB, RM	Mulching, tumbleweed, HWY I-40, grasses on other side of HWY.
16.12 _1_NE SW	12/8/2 021 8:59am	100- 0380	35.12199 482	- 107.82351 06	W	Tumbleweed /mulching @23' @270°	Blue exit sign @827' @270°	Olympus red	AM, CB, RM	Mulching, tumbleweed, fence, HWY I-40, bridge, and mountain range.
16.12 _2_NE SW	12/8/2 021 10:10a m	100- 0392	35.13109 979	- 107.82293 55	N	Fourwing salt bush @18' @360°	Little hil @134' @360°	Olympus red	AM, CB, RM	Mountain range way in the back

16.12 _2_NE SW	12/8/2 021 10:10a m	100- 0393	35.13109 979	- 107.82293 55	E	Salt cedar @133' @90°	Hill @204' @90°	Olympus red	AM, CB, RM	Grassy, dirt road, drainage ditch, and mountain range.
16.12 _2_NE SW	12/8/2 021 10:10a m	100- 0396	35.13109 979	- 107.82293 55	S	Grass patch @1-22' @180°	Salt cedar @935' @180°	Olympus red	AM, CB, RM	Grassy, playa/salt, salt cedar, and mountain range.
16.12 _2_NE SW	12/8/2 021 10:10a m	100- 0398	35.13109 979	- 107.82293 55	W	Grassy patch/tire @12' @270°	Quality Inn Motel @837' @270°	Olympus red	AM, CB, RM	Grassy, salt cedar, motel, and mountain range.

Appendix II - Photos



2016: 16.12_1_N.
North view inside of
polygon 1.

2021: 16.12_1_N



2016:
16.12_1_E. East
view inside of
polygon 1.



2021:
16.12_1_E



2016: 16.12_1_S.
South view inside of
polygon 1.

2021: 16.12_1_s



2016:
16.12_1_W.
West view
inside of



2021:
16.12_1_
W



2016:
16.12_2_N
. North
view from
polygon 1,
looking
into
polygon 3.

2021:
16.12_2_
N



2016:
16.12_2_E.
East view
inside of
polygon 1.



2021: 16.12_2_E

2016:
16.12_2_S.
South view
inside of
polygon 1.



2021: 16.12_2_S



2016:
16.12_2_W.
West view
inside of
polygon 1.

2021:
16.12_2_W



Appendix III – Current monitoring methods available

Low-intensity methods

- Where: happens on all sites with GRGWA projects
- Method name: NMRAM (New Mexico Rapid Assessment Method v 2.1)
- Time required: 3 hours – half day/ site
- Repeat: done once pre-treatment and in 4-5 year intervals post-treatment
- Basics: mapping vegetation communities (by vertical and horizontal structure), recording dominant vegetation in each strata (trees, shrubs, herbaceous), assessing fuel load, noting soil surface condition and native/exotic ratio at all vegetation levels, photo points
- Any on-site impacts or materials: none

High-intensity methods

- Where: happens on select sites, in addition to low-intensity monitoring

Submethod name 1: BBIRD or BEMP vegetation plots (depends on treatment area size)

- Time required: approx. 2 hours/site
- Repeat: both pre-treatment and in 4-5 yr intervals post-treatment
- Basics: larger plots and transects documenting vegetation, photo points
- On-site impacts or materials: rebar and cap

Submethod name 2: Brown's transects

- Time required: 1-1.5 hours/site
- Repeat: both pre-treatment and in 4-5 yr intervals post-treatment
- Basics: transects to calculate fuel loading and fire behavior, photo points
- On-site impacts or materials: rebar and cap

Submethod name 3: BEMP-adapted Groundwater Well Monitoring

- Time required:
 - Initial installation: 1-2 hours/ well (ideally 2+ wells/site)
 - Repeat: maintenance as needed, should be minimal
 - Data offloading: 10-20 minutes/well
 - Repeat: at least annually (this is when we anticipate datalogger will be full and batteries will need to be changed)
- Basics: install a well with a sensor which records groundwater level and temperature once an hour year round; this will reflect changes due to seasonal variation, vegetation growth, irrigation, etc.
- On-site impacts or materials: shallow monitoring well (consists of capped PVC pipe extending into the ground about 3 feet below the water table and above ground approx. 2 feet (can be painted earth tones); well contains a datalogger (pressure transducer) suspended on a cable into the water); well should be protected from cattle grazing (so may require rebar around pvc visible above ground)

Appendix IV - Modified Hink and Ohmart categories, from NMRAM

The following is pages 39-41 in Muldavin et al.'s 2014 NMRAM for Montane Riverine Wetlands v 2.0 Manual (draft, not yet published)

Vegetation Vertical Structure Type Definitions for NMRAM

Multiple-Story Communities (Woodlands/Forests)



Type 1 – High Structure Forest with a well-developed understory.

Tall mature to intermediate-aged trees (>5 m [>15 feet]) with canopy covering >25% of the area of the community (polygon) and understory layer (0-5 m [0-15 feet]) covering >25% of the area of the community (polygon). Substantial foliage is in all height layers. (This type incorporates Hink and Ohmart structure types 1 and 3.) Photograph on Gila River by Y. Chauvin, 2012.



Type 2 –Low Structure Forest with little or no understory.


Tall mature to intermediate-aged trees (>5 m [>15 feet]) with canopy covering >25% of the area of the community (polygon) and understory layer (1-5 m [3-15 feet]) covering <25% of the area of the community (polygon). Majority of foliage is over 5 m (15 feet) above the ground. (This type incorporates Hink and Ohmart structure types 2 and 4.) Photograph on Diamond Creek by Y. Chauvin, 2012.

Single-story Communities (Shrublands, Herbaceous and Bare Ground)



Type 5 –Tall Shrub Stands.


Young tree and shrub layer only (15-5 m [4.5-15 feet]) covering >25% of the area of the community (polygon). Stands dominated by tall shrubs and young trees, may include herbaceous vegetation underneath the woody vegetation. Photograph on San Francisco River by Y. Chauvin, 2012.

**Type 6S-Short Shrub Stands.**

Short stature shrubs or very young shrubs and trees (up to 1.5 m [up to 4.5 feet]) covering >10% of the area of the community (polygon). Stands dominated by short woody vegetation, may include herbaceous vegetation underneath the woody vegetation. Photograph on Lower Pecos River by E. Lindahl, 2008.

**Type 6W-Herbaceous Wetland.**

Herbaceous wetland vegetation covering >10% of the area of the community (polygon). Stands dominated by obligate wetland herbaceous species. Woody species absent, or <10% cover. Photograph of *Carex nebrascensis* meadow on upper Rio Santa Barbara by Y. Chauvin, 2009.

**Type 6H- Herbaceous.**

Herbaceous vegetation covering >10% of the area of the community (polygon). Stands dominated by herbaceous vegetation of any type except obligate wetland species. Woody species absent or <10% cover. Photograph on Diamond Creek by Y. Chauvin, 2012.



Type 7–Sparse Vegetation/Bare Ground.

Bare ground, may include sparse woody or herbaceous vegetation, but total vegetation cover <10%. May be natural in origin (cobble bars) or anthropogenic in origin (graded or plowed earth) Photograph on Lower Gila River by Y. Chauvin, 2012.