

Ojo Caliente Project 16-16

Pre-treatment Monitoring Report

2016



Prepared by

Kathryn R Mahan, Ecological Monitoring Specialist & Daniel Hernandez, Ecological Monitoring Technician, New Mexico Forest and Watershed Restoration Institute for the Greater Rio Grande Watershed Alliance



Contents

| | |
|---|----|
| Acronyms and Abbreviations | 3 |
| Purpose of Report | 4 |
| Ecological Context of Bosque Restoration | 4 |
| Monitoring and Field Methods | 5 |
| Low intensity Field Methods | 5 |
| High-intensity Field Methods | 6 |
| Estimating Vegetation Cover using eCognition Software | 6 |
| Personnel Involved | 7 |
| Ojo Caliente Project | 7 |
| Discussion | 21 |
| References | 22 |
| Appendix I - Photopoint Table | 24 |
| Appendix II - Photos | 25 |
| Appendix III – Current monitoring methods available | 32 |
| Appendix IV - Modified Hink and Ohmart categories, from NMRAM | 33 |

Acronyms and Abbreviations

| Acronym, Abbreviation, or Term | Explanation or Definition as used by NMFWR I |
|---------------------------------------|--|
| AGL | above ground level; GIS term |
| BBIRD plots | Breeding Biology Research and Monitoring Database, larger circular plot types |
| BEMP plots | Bosque Ecosystem Monitoring Program, small rectangular plot types |
| FEAT | Fire Ecology Assessment Tool |
| FFI | FEAT/ FIREMON Integrated |
| FIREMON | Fire Effects Monitoring and Inventory System |
| FSA | Farm Service Agency, a department of the USDA |
| GIS | Geographic Information Systems |
| GRGWA | Greater Rio Grande Watershed Alliance |
| LIDAR | Light detecting and ranging, a remote sensing technique using light to gather elevation data |
| NAIP | National Agriculture Imagery Program (aerial imagery) |
| NDVI | Normalized Difference Vegetation Index; GIS term for a band ratio of the visible red and the near infrared spectral bands and is calculated using the following formula: $(NIR - Red)/(NIR + Red)$ |
| NHNM | Natural Heritage New Mexico |
| NMDGF | New Mexico Department of Game and Fish |
| NMED SWQB | New Mexico Environment Department Surface Water Quality Bureau |
| NMFWR I | New Mexico Forest and Watershed Restoration Institute |
| NMHU | New Mexico Highlands University |
| NMRAM | New Mexico Rapid Assessment Method, version 2.0 |
| NRCS | Natural Resource Conservation Service |
| PC | Plot center |
| RGIS | Resource Geographic Information System |
| SWCD | Soil and Water Conservation District |
| TIFF | Tagged image file format |
| USDA | United States Department of Agriculture |
| USGS | United States Geological Survey |
| WQCC | Water Quality Control Commission |
| WSS | Web Soil Survey, a soils database of the NRCS |

Purpose of Report

This report covers the low-intensity pre-treatment vegetation monitoring assessment performed on a non-native phreatophyte removal project submitted for the Ojo Caliente River to the Greater Rio Grande Watershed Alliance in 2016. Following a discussion of the ecological context, and our monitoring methods, we present pertinent background, observations, and assessment results for the project.

Ecological Context of Bosque Restoration

Neither the challenges nor the importance of working in the bosque and other riparian areas in New Mexico today should be underestimated. According to the New Mexico Department of Game and Fish Conservation Division, wetlands and riparian areas comprise approximately 0.6 percent of all land in New Mexico (2012). Despite this small percentage, estimates of New Mexican vertebrate species depending on wetland and riparian habitat for their survival ranges from 55% (New Mexico Department of Game and Fish Conservation Services Division, 2012) to 80% (Audubon New Mexico, 2013). These areas also provide flood mitigation, filtration of sediment and pollutants, and water for a variety of purposes including groundwater recharge (Audubon New Mexico, 2013). In addition, native vegetation such as cottonwoods have cultural significance to many communities.

As much as these areas are disproportionately important to ecosystems and human communities, they are equally disproportionately impacted by disturbance. Anthropogenic impacts with major consequences for our riparian areas include dams, reservoirs, levees, channelization, acequias and ditches, jetty jacks, riprap and Gabion baskets, urbanization, removal of native phreatophytes, grazing by domestic livestock, excessive grazing pressure by native ungulate populations absent natural predation cycles, beaver removal, logging, mining, recreation, transportation, introduction and spread of invasive exotic species, groundwater extraction, altered fire and flood regimes drought and climate change (Committee on Riparian Zone Functioning and Strategies for Management, et al., 2002). Statewide, it is estimated that as much as 90% of New Mexico's historical riparian areas have been lost (Audubon New Mexico, 2013), and approximately 39% of our remaining perennial stream miles are impaired (New Mexico Department of Game and Fish Conservation Services Division, 2012).

New Mexico is fortunate enough to have the Middle Rio Grande Bosque, the largest remaining bosque in the Southwest (USDA USFS, 1996). However, over the past two decades, the number of fires in the bosque has been increasing. Historically, the primary disturbance regime in the bosque has been flooding, not fire, which means the system is not fire-adapted. In fact, native species like cottonwood resprout from their roots after floods and need wet soils to germinate from seed. Flooding also promotes decomposition of organic material and keeps the soil moist which reduces the likelihood of fire. Today, overbank flow is uncommon in many areas of the Rio Grande due to the heavy alteration of the channel and flow regimes (two obvious examples are the structures defining the upper and lower extent of the Middle Rio Grande: Cochiti Dam and Elephant Butte Reservoir). This has led to low fuel moisture content and high fuel loads, as well as increased human presence in the riparian area. As a result, bosque fires are more common and more severe: they kill cottonwoods and other native species, creating spaces which are filled by non-native species such as salt cedar, Russian olive, Siberian elm, and Tree-of-Heaven. We are constantly learning more about how these species can exploit and encourage a riparian fire regime, in addition to many other changes they bring to ecosystems.

Efforts geared toward the removal of these nonnative species can help to reduce fire risk, preserve native vegetation, and be part of a larger effort to restore the bosque and the watershed as a whole to a more natural and functional ecosystem. The Greater Rio Grande Watershed Alliance (GRGWA) has been working on these issues with a variety of collaborating organizations and agencies within the Rio Grande basin for several years. Since 2013, the New Mexico Forest and Watershed Restoration Institute (NMFWR) has been working with GRGWA and the Claunch-Pinto Soil and Water Conservation District (SWCD) to begin construction of a geodatabase for all of GRGWA's non-native phreatophyte removal projects as well as to perform the formal pre- and post-treatment monitoring, utilizing the field methods explained below as well as LIDAR analysis where appropriate and available.

Monitoring and Field Methods

Low intensity Field Methods

Low intensity pre-treatment vegetation monitoring was done using an adapted version of the biotic portion of the New Mexico Rapid Assessment Method (NMRAM), v 2.1, updating recommendations made in the Field Manual for Greater Rio Grande Watershed Alliance (GRGWA) Riparian Restoration Effectiveness Monitoring and the GRGWA Monitoring Plan, developed by Lightfoot & Stropki of SWCA Environmental Consultants in 2012. (For a brief overview of both low and high intensity monitoring methods used by the NMFWR on GRGWA projects, please see Appendix III.)

For those not familiar, NMRAM was developed by the New Mexico Environment Department Surface Water Quality Bureau Wetlands Program and Natural Heritage New Mexico as a "cost effective, yet consistent and meaningful tool" (Muldavin, 2011) for wetland ecological condition assessment in terms of anthropogenic disturbance as negatively correlated with quality and functionality. The portions of NMRAM we utilized are Level 2 "semi-quantitative" field measurements taken at less detail than plot level (Muldavin, 2011).

Measurements taken included relative native plant community composition, vegetation horizontal patch structure, vegetation vertical structure, native riparian tree regeneration, and invasive exotic plant species cover. The underlying method for these biotic assessments was a version of the 1984 Hink and Ohmart vertical structure classification system, modified for use in the NMRAM for Montane Riverine Wetlands version 2.0 (see Appendix IV). First, vegetation communities were mapped out by patch (polyon) according to the Hink and Ohmart system. Next, the presence of (state-listed) invasives, wetland species, and the two dominant species in each strata ("tree" >15 ft, "shrub" 4.5-15 ft, and "herbaceous" <4.5 ft) were recorded for each plant community. The native/exotic ratio in each of the patches was scored and weighted based on the percent of the project area each patch comprised. These scores were then combined with the additional biotic metrics of vertical and horizontal diversity, native tree regeneration, and overall (listed) invasive presence. The NMRAM rating system is based, on all levels, on a scale of 1 to 4, where 4 is considered excellent condition, 3 good, 2 fair, and 1 poor.


We also assessed soil surface condition, which is a metric typically included in the abiotic section of the NMRAM, as well as the presence of surface fuels, which is not part of the NMRAM. Unlike the other 6 metrics we used, surface fuels were recorded on a rating scale from 0 to 1.0 where 1.0 is a continuous fuel matrix.

Photopoints were established to capture images where vegetation shifts were observed and/or at representative locations throughout the site. Waypoints were marked with a Garmin GPS unit and named sequentially by site. Photos were taken facing north, east, south and west at each point.

Prior to entering the field, we created a map with the project boundaries as provided by GRGWA. We combined these polygons with recent aerial imagery and identified relevant roads and other landscape features. Once on the ground, the vegetation community polygons (as determined by the modified Hink and Ohmart classification system) were hand-drawn onto this map and served as the basis for other biotic metric assessments. Upon return to the office, this polygon map and the photopoints were digitized by the monitoring technician and/or specialist.

High-intensity Field Methods

High-intensity monitoring was also done, in part, on this site. We used an adapted Bosque Ecosystem Monitoring Program (BEMP) style plot. These are 16 x 98-foot rectangles, placed approximately parallel to the river. Within these plots, we measure canopy and species, and vegetation and ground cover. We also used Brown's transects to measure surface fuels.



Estimating Vegetation Cover using eCognition Software

Object based image classification systems, such as eCognition software, allows for a semi-automated analysis of high resolution images. This approach divides the image into meaningful homogenous regions, known as image objects. These image objects are groups of pixels that are adjacent to each other and are spectrally similar. Once image objects are created, they provide a great deal of information from which an image classification can be developed (Lizarazo & Elsner, 2009). In large areas where more detailed vegetation surveys are cost prohibited, eCognition provides a means to characterize a landscape using readily available aerial photography.

For the Ojo Caliente Area LIDAR, light detecting and ranging elevation data, was not available. Instead, digital ortho-imagery was used to estimate vegetation areas. 2014 NAIP (National Agriculture Imagery Program) imagery was acquired for the study area. NAIP is a USDA/FSA program to acquire 'leaf on' aerial imagery during the peak growing season. NAIP imagery for New Mexico can be downloaded by Quarter Quadrangle extent in an uncompressed TIFF format via RGIS –Resource Geographic Information System (<http://rgis.unm.edu/>).

NAIP was collected in 2014 with the near infrared (NIR) spectral band. The 4 band imagery (Red, Green, Blue, and NIR) at 1 meter cell size is available statewide. Having the NIR band allows for a greater analysis of vegetation and the calculation of the Normalized Difference Vegetation Index (NDVI). NDVI is a band ratio of the visible red and the near infrared spectral bands and is calculated using the following

formula: $(NIR - Red)/(NIR + Red)$. This makes vegetation monitoring and analysis feasible over large areas.

The classification was based on finding the right threshold values for each feature. To determine specific threshold values, information about each image object could be displayed and tested to determine if those values were appropriate for the given land cover feature.

To identify dominant vegetation types, the image was first classified to three classes: Vegetation, Bare Ground, and Water. Image properties of Hue, Saturation, and Intensity and NDVI values of the image objects were used to identify Vegetation and Bare Ground (Type 7) classes. Large negative near infra-red values were used to identify water features. The vegetation classification was further stratified to separate forested areas from herbaceous vegetation using image texture features. Texture features are created in eCognition to determine the arrangement and frequency of tonal variation in certain areas of an image. To give texture values, eCognition uses differences in grey level differences and contrast. For example, areas of forests and shrub have a higher texture value than smooth flat grassland areas. After herbaceous vegetation (Type 6H) was separated from Forest/Shrub vegetation, existing field data was used to divide the Forest/Shrub vegetation into Type 1 (High structure forest with understory) and Type 5 (Tall Shrubs) vertical structure type, and Type 6S (short shrubs). The classification was exported from eCognition and then manually edited using Erdas Imagine software. Vertical structure types were interpreted using the imagery, photographs and field data collected on site. The resulting classification was used to determine acreage totals by vertical structure type according to the NMRAM definitions.

Personnel Involved

2016 New Mexico Forest and Watershed Restoration Institute Monitoring Team:

- Kathryn R Mahan, Ecological Monitoring Specialist
- Dr Rob Strahan, Restoration Monitoring Program Manager
- Daniel Hernandez, Ecological Monitoring Technician

2016 New Mexico Forest and Watershed Restoration Institute GIS Team:

- Patti Dappen, GIS Specialist

Other persons contacted:

- Fred Rossbach, Field Coordinator, Greater Rio Grande Watershed Alliance
- Peter Vigil, Taos Soil and Water Conservation District

Ojo Caliente Project

Project 16-16 is located Rio Arriba County in Ojo Caliente, N.M, along the Rio Ojo Caliente and in fenced pastures west of US 285. It is approximately 25 miles north of Española.

The average precipitation for the nearby city of Española is 11.41 inches per year. The average high temperature is 90° F in July and the average low is 15° F in December and January (U.S. Climate Data, 2017). According to the NRCS Web Soil Survey, the project area is comprised of 89% Fluvents, nearly level, 9.4% Royosa loamy sand, 1 to 8 percent slopes, 1.3% Sedillo-Orthents association, strongly sloping.

Ecological sites within this project include R035XA112NM Loamy, R035XA113NM Sandy, R035XG114NM Gravelly, R036XA004NM Gravelly Slopes, R036XB006NM Loamy, R036XB007NM Malpais, R036XB008NM Meadow, R036XB011NM Sandy, R036XB111NM Sandy Slopes, and R036XB132NM Gravelly Hills (USDA NRCS, 2016).

The Loamy ecological site (R035XA112NM) typically supports a grassland state dominated by blue grama, western wheatgrass, galleta, ring muhly, dropseeds, and/or threeawns. It can also be found in a piñon-juniper invaded state (dominated by piñon, juniper, and blue grama), a grass/succulent-mix state (dominated by blue grama, cholla and prickly pear), a shrub-dominated state (dominated by rabbitbrush or horsebrush and blue grama), as well as a bare state with sparse grass (USDA NRCS n.d.).

The Sandy ecological site typically supports plant communities composed of fourwing saltbush, winterfat, and sagebrush at the shrub layer, and at the herbaceous layer, Rocky Mountain bee plant, blue grama, western wheatgrass, threeawns, galleta, dropseed, Indian ricegrass, needle-and-thread, squirreltail, and New Mexico feathergrass. This may also support a shrub-dominated state (dominated by sagebrush, rabbitbrush with a blue-grama/threeawn/dropseed/muhly understory), as well as a juniper-dominated state (with a patchy grass understory of blue grama, dropseeds, galleta, Indian ricegrass and threeawn) (USDA NRCS n.d.).

The Gravelly ecological site type typically supports grassland with minor shrub and piñon-juniper components. Common dominant grass species include blue, black and sideoats grama, little bluestem, spike muhly, Western wheatgrass, New Mexico feathergrass, Indian ricegrass, and squirreltail. Common shrubs include fourwing saltbush, winterfat, Apache plume, rabbitbrush, soapweed yucca, and sagebrush and broom snakeweed. The site can also be found in a shrub-encroached state dominated by rabbitbrush and blue grama; erosion is more common in this state (USDA NRCS n.d.).

The Gravelly slopes ecological site is usually mixed grasses and shrubs including needlegrass, Indian ricegrass, western wheatgrass, sideoats grama, galleta, blue grama, dropseeds, threeawn fourwing saltbush and winterfat. Piñon/juniper trees may also be present in small amounts. Continuous grazing has been observed to shift the species balance to less desirable grasses and woodies (USDA NRCS n.d.).

The Loamy ecological site (R036XB006NM) is a grassland site with scattered shrubs throughout. Forbs are a minor component. Other species include sideoats grama, sand dropseed, pine dropseed, mat muhly, cheatgrass, pingue, woolly Indianwheat, globemallow, prairie coneflower, oneseed juniper, piñon pine, pale wolfberry, broom snakeweed, yucca, cholla cactus and antelope bitterbrush (USDA NRCS n.d.).

The Malpais ecological site typically supports a grassland state dominated by blue grama and sideoats grama. Other common vegetation includes western wheatgrass, little bluestem, spike muhly, black grama, galleta, New Mexico feathergrass, alkali sacaton, winterfat, fourwing saltbush, broom snakeweed, and scattered piñon and juniper. In a deteriorated state, the grass community may become sod-bound, dominated by blue grama sod, threeawns, wolfstail and snakeweed (USDA NRCS n.d.).

The Meadow ecological site type contains approximately 90 to 95 percent vegetation suitable for grazing or browsing. However, due to the high availability of soil moisture, which results in early green up and high productivity, this site is subject to deterioration by overgrazing and trampling. Deterioration

is indicated by a decrease in western wheatgrass, tufted hairgrass, bromes, and bluegrass with an increase in mat muhly, sedges, rushes, and forbs (USDA NRCS n.d.).


The Sandy ecological site type is typically mixed-grass and shrub. Fourwing saltbush and winterfat are the dominant shrubs with big sagebrush and rabbitbrush occurring in lesser amounts. Few, if any, trees occur on this site. Forbs are a minor component except during spring (USDA NRCS n.d.).

The Sandy Slopes ecological site typically supports a mixed-shrub grassland state dominated by sagebrush, saltbush, winterfat, rabbitbrush and sagewort at the shrub level, and blue grama, galleta, dropseed, Indian ricegrass, and threeawn at the herbaceous level. It can also be found in Piñon-juniper state (up to 15% piñon-juniper cover with patchy grass), shrub state (sagebrush, saltbush and squirreltail), and an eroded state (piñon-juniper, annual forbs/grasses, bare ground and reduced A-horizon in the soil) (USDA NRCS n.d.).

The Gravelly Hills ecological site typical plant community consists of sideoats and black grama grasses. Shrubs include Bigelow sagebrush, broom snakeweed, longleaf ephedra, feather dalea and yucca. Sparse strands of piñon-juniper may be widely scattered across the site. Overgrazing may cause loss of grass cover allowing piñon and juniper seedlings to establish and possibly facilitate the transition to piñon/juniper encroachment (USDA NRCS n.d.).

Monitoring was conducted at this 143.8-acre site on November 11, 2016 as part of an initial treatment as part of a restoration project targeting non-native phreatophytes scheduled for 2016-2017. The project is located in Ojo Caliente, NM in Rio Arriba County along the Rio Ojo Caliente. The Taos Soil and Water Conservation District sponsored the project. Portions of the project are accessed through the Ojo Caliente Spa. The project is an initial treatment to remove salt cedar, Russian olive, and Siberian elm trees by extraction and mastication, although very large trees will not be treated, and enough trees will be left to create 10-20% canopy cover. Restoration goals include returning the riparian area to a more natural state, promoting native plant species, enhancing a wildlife habitat corridor, reducing fire hazard and continuing to long-term watershed health and ecosystem function.

16-16 Ojo Caliente



16-16 Ojo Caliente is located in Ojo Caliente, N.M. in Rio Arriba County along the Rio Ojo Caliente. Portions of the project are accessed through the Ojo Caliente Spa. The project is an initial treatment to remove salt cedar, Russian olive, and Siberian elm.



Figure 1. Project 16-16 in geographic context.

Ojo Caliente - Taos SWCD - 16-16



Figure 2. 16-16 Ojo Caliente project outline.

The Rio Ojo Caliente, as visible in the image above, can take different paths depending on the amount of flow. The Rio Ojo Caliente flows from the Tusas and Vallecitos near Lamadera through the Ojo Caliente valley, to join the Rio Chama near Chili, south of Medenales and north of El Duende and Hernandez. The USGS gauge 08289000 measures the Rio Ojo Caliente at Lamadera.

The GRGWA site is located approximately 6 miles south of the gauge at an elevation around 6200 feet. Exotic species observed included Russian olive, Siberian elm, salt cedar, tree-of-heaven, cheatgrass, sweetclover, Russian thistle, and musk thistle. Native plants observed at the time of the site visit included Rio Grande cottonwood, oneseed juniper, oak, seepwillow, New Mexico olive, coyote willow, rubber rabbitbrush, broom snakeweed, sumac, cattail, wild licorice, alkali sacaton, dropseed and other grasses.

Table 1. NMRAM Scores for 16-16.

| Metric 16.16 November 4, 2016 | Score |
|---|--------------|
| Relative Native Plant Community Composition | 1 |
| Vegetation Horizontal Patch Structure | 4 |
| Vegetation Vertical Structure | 3 |
| Native Riparian Tree Regeneration | 2 |
| Exotic Invasive Plant Species Cover | 1 |
| | |
| Project Biotic Score (based on above ratings) | 2.2 |
| Project Biotic Rating | C/Fair |
| | |
| Soil Surface Condition | 1 |
| Surface Fuels | 0.2 |

The lowest scores for this project came in the Relative Native Plant Community Composition and Exotic Invasive Plant Species Cover metrics, due to the high percentage of invasive plants. The project scored best in the vegetation horizontal and vertical structure metrics, because there are several different plant communities distributed across the landscape and a dominance of overstory structure. Vegetation polygons are represented by structure type in the map, Figure 3 and Figure 4, as well as in the NAIP vegetation classification map, Figure 5 through Figure 7. Overall, this site scored a 2.1 out of 4 overall, which is a “C” or “Fair” biotic rating.

This site also had one plot established (location shown on map below). At this plot, we collected data on vegetation cover and fuel loading using Submethods 1 and 2 outlined in Appendix III, the BEMP plots and the Brown’s transects.

16-16 North Ojo Caliente Taos SWCD




Figure 3. 16-16 Ojo Caliente project vegetation polygons – northern portion.

16-16 South Ojo Caliente Taos SWCD




Figure 4. 16-16 Ojo Caliente project vegetation polygons – southern portion.

16-16 Ojo Caliente (North) - Vertical Structure Classification




Figure 5. NAIP vegetation classification for 16-16 – North third only.

16-16 Ojo Caliente (Middle) - Vertical Structure Classification




Figure 6. NAIP vegetation classification for 16-16 – Middle third only.

16-16 Ojo Caliente (South) - Vertical Structure Classification




Figure 7. NAIP vegetation classification for 16-16 – Southern third only.

Table 2. 16-16 Average surface fuels from 2 transects on plot.

| Fuel | Average tons/acre |
|-----------------|--------------------|
| 1-hr | 0.04 |
| 10-hr | 0.03 |
| 100-hr | 0.01 |
| 1000-hr | 11 |
| All woody fuels | 11 |
| | |
| Fuel | Avg depth (inches) |
| Duff | 0.38 |
| Litter | 2.3 |
| Total | 2.7 |




Figure 8. Percent Vegetative Cover for plot on 16-16.




Figure 9. Percent Vegetative Cover for plot on 16-16.




Figure 10. Percent ground cover for plot on 16-16.




Figure 11. Percent ground cover for plot on 16-16.

Discussion

We would like to clarify that we are adapting these NMRAM metrics for our own purposes. That is, we are using them both inside and outside their intended site ranges, including on larger sites (NMRAM is designed to handle a site around 100 x 200 meters), sites further from the river (NMRAM is currently in use primarily for assessing riverine wetlands), and sites defined by exotic vegetation presence rather than hydrologic boundaries and upland vegetation indicators/apparent wetland extent. Site delineation and size is likely to be variable for a number of other reasons, including landowner participation, available funds, proposals received from contractors, etc – many of which cannot be directly correlated to site disturbance or ecological function. For this reason, we do not use the entire NMRAM assessment, or place confidence in the weighted score roll-ups that are typically part of an NMRAM report. Should one be interested, rationale for the weighting in the NMRAM score roll-up can be found in the yet-to-be-published field manual for version 2.1. For more information, contact Maryann McGraw of the NMED or NMFWR.

While we provide a biotic site score and rating for your reference, we recommend comparisons be done with individual metrics from pre-treatment and post-treatment assessment from the same site, rather than across multiple sites. Also of note is that statistical analysis is not appropriate for NMRAM, or other low intensity, rapid field methods.

Please note that should the project area change significantly from what was originally proposed and monitored, all metrics will lose some amount of confidence on comparison as it is impractical to re-examine the original site assessment scores using new boundaries. This is an issue of concern of which GRGWA should be aware. We recommend that GRGWA attempt to minimize alterations in project boundaries once pre-treatment monitoring data has been approved for collection. Another, somewhat alternative, recommendation is that the initial monitoring regime include high-intensity modified BEMP-type plots which could be repeated in their exact initial locations, allowing collection of comparable data regardless of boundary change. We recognize that this is not always practical: boundaries change for a number of reasons and time and cost constraints can necessitate the sole use of a rapid assessment method for monitoring. We have reason to hope our outlined assessment method will still be a satisfactory indicator for site function improvement or degradation primarily because metrics in rapid assessment methods such as this are set up to have relatively low sensitivities (i.e. for a change to be reflected in the metrics, either positive or negative, disturbance on site has to be significantly altered).

From here on out, the goal of the GRGWA/ NMFWR is that all sites will be revisited for post-treatment monitoring in 5-year intervals. It is our intention and expectation that the data collected in these intervals will reflect any significant changes in disturbance and ecological function of the site.

References

- Audubon New Mexico. (2013). *Water Matters: Water for New Mexico Rivers*. Albuquerque, New Mexico: Utton Transboundary Resources Center.
- Brown, J. K. (1974). *Handbook for Inventorying Downed Woody Material*, USDA Forest Service General Technical report INT-16. *Handbook for Inventorying Downed Woody Material*. Ogden, Utah: USDA Forest Service Intermountain Forest and Range Experiment Station.
- Bureau of Land Management. (2006). *Grazing Management Processes and Strategies for Riparian-Wetland Areas, TR 1737-20*.
- Claunch-Pinto Soil and Water Conservation District on behalf of the Greater Rio Grande Watershed Alliance. (2015). *Request for Proposals for Greater Rio Grande Watershed Alliance Riparian Restoration Projects*. Mountainair, NM: Claunch-Pinto Soil and Water Conservation District.
- Committee on Riparian Zone Functioning and Strategies for Management, et al. (2002). *Riparian Areas: Functions and Strategies for Management*. Washington, D.C.: National Academy Press.
- Lightfoot, D. &. (2012). *Greater Rio Grande Watershed Alliance Riparian Restoration Effectiveness Monitoring Plan*. Albuquerque, NM: SWCA Environmental Consultants.
- Lightfoot, David & Stropki, C. (2012). *Field Manual for Greater Rio Grande Watershed Alliance Riparian Restoration Effectiveness Monitoring*. Albuquerque, NM: SWCA Environmental Consultants.
- Lizarazo, I., & Elsner, P. (2009). Fuzzy segmentation for object-based image classification. *International Journal of Remote Sensing*, 30.
- Muldavin, E. B. (2011). *New Mexico Rapid Assessment Method: Montaine Riverine Wetlands*. Version 1.1. Final report to the New Mexico Environment Department, Surface Water Quality Bureau. 90 pp. and appendices.
- New Mexico Department of Game and Fish Conservation Services Division. (2012). *Bridge and Road Construction/Reconstruction Guidelines for Wetland and Riparian Areas*.
- U.S. Climate Data. (2017). *Climate New Mexico*. Retrieved from U.S. Climate Data: <http://www.usclimatedata.com/climate/new-mexico/united-states/3201>
- USDA NRCS. (2013, December 6). *Web Soil Survey*. Retrieved from <http://websoilsurvey.sc.egov.usda.gov/App/HomePage.htm>
- USDA NRCS. (2016, 8 10). *Web soil Survey*. Retrieved from <https://websoilsurvey.sc.egov.usda.gov/App/HomePage.htm>
- USDA NRCS n.d. (n.d.). *Ecological Site Description Gravelly R035XG114NM*.
- USDA NRCS n.d. (n.d.). *Ecological Site Description Loamy R036XB006NM*.
- USDA NRCS n.d. (n.d.). *Ecological Site Description Gravelly Hills R036XB132NM*.
- USDA NRCS n.d. (n.d.). *Ecological Site Description Gravelly Slopes R036XA004NM*.
- USDA NRCS n.d. (n.d.). *Ecological Site Description Loamy R035XA112NM*.

USDA NRCS n.d. (n.d.). *Ecological Site Description Malpais R036XB007NM* .

USDA NRCS n.d. (n.d.). *Ecological Site Description Meadow R036XB008NM* .

USDA NRCS n.d. (n.d.). *Ecological Site Description Sandy R035XA113NM* .

USDA NRCS n.d. (n.d.). *Ecological Site Description Sandy R036XB011NM* .

USDA NRCS n.d. (n.d.). *Ecological Site Description Sandy Slopes R036XB111NM* .

USDA USFS. (1996, September). Ecology, Diversity, and Sustainability of the Middle Rio Grande Basin, RM-GTR-268. (D. M. Finch, & J. A. Tainter, Eds.) Fort Collins, Colorado.

Appendix I - Photopoint Table

| Name | Latitude | Longitude |
|---------------|-----------------|------------------|
| 16.16_1_NESW | 36.3085 | -106.0460 |
| 16.16_2_NESW | 36.3104 | -106.0450 |
| 16.16_3_NESW | 36.3212 | -106.0390 |
| 16.16_4_NESW | 36.3009 | -106.0530 |
| 16.16_PC_NESW | 36.3194 | -106.0410 |

Appendix II - Photos



16.16_1_N. View facing north inside of polygon 2.

16.16_1_E. View facing east inside of polygon 2.



16.16_1_S. View facing south inside of polygon 2.

16.16_1_W. View facing west from inside polygon 2.



16.16_2_N. View facing north from inside polygon 4.

16.16_2_E. View facing east from inside of polygon 4.





16.16_2_S. View facing south from inside of polygon 4.

16.16_2_W. View facing west inside of polygon 4.



16.16_3_N. View facing north inside of polygon 6.

16.16_3_E. View facing east from inside polygon 6.



16.16_3_S. View facing south from inside polygon 6.

16.16_3_W. View facing west from inside of polygon 6.





16.16_4_N. View facing north from inside of polygon 7.

16.16_4_E. View facing east inside of polygon 7.



16.16_4_S. View facing south inside of polygon 7.

16.16_4_W. View facing west from inside polygon 7.



16-16 Plot Photos



N from PC



E from PC



S from PC



W from PC



Plot Center

Appendix III – Current monitoring methods available

Low-intensity methods

- Where: happens on all sites with GRGWA projects
- Method name: NMRAM (New Mexico Rapid Assessment Method v 2.1)
- Time required: 3 hours – half day/ site
- Repeat: done once pre-treatment and in 4-5 year intervals post-treatment
- Basics: mapping vegetation communities (by vertical and horizontal structure), recording dominant vegetation in each strata (trees, shrubs, herbaceous), assessing fuel load, noting soil surface condition and native/exotic ratio at all vegetation levels, photo points
- Any on-site impacts or materials: none

High-intensity methods

- Where: happens on select sites, in addition to low-intensity monitoring

Submethod name 1: BBIRD or BEMP vegetation plots (depends on treatment area size)

- Time required: approx. 2 hours/site
- Repeat: both pre-treatment and in 4-5 yr intervals post-treatment
- Basics: larger plots and transects documenting vegetation, photo points
- On-site impacts or materials: rebar and cap

Submethod name 2: Brown's transects

- Time required: 1-1.5 hours/site
- Repeat: both pre-treatment and in 4-5 yr intervals post-treatment
- Basics: transects to calculate fuel loading and fire behavior, photo points
- On-site impacts or materials: rebar and cap

Submethod name 3: BEMP-adapted Groundwater Well Monitoring

- Time required:
 - Initial installation: 1-2 hours/ well (ideally 2+ wells/site)
 - Repeat: maintenance as needed, should be minimal
 - Data offloading: 10-20 minutes/well
 - Repeat: at least annually (this is when we anticipate datalogger will be full and batteries will need to be changed)
- Basics: install a well with a sensor which records groundwater level and temperature once an hour year round; this will reflect changes due to seasonal variation, vegetation growth, irrigation, etc.
- On-site impacts or materials: shallow monitoring well (consists of capped PVC pipe extending into the ground about 3 feet below the water table and above ground approx. 2 feet (can be painted earth tones); well contains a datalogger (pressure transducer) suspended on a cable into the water); well should be protected from cattle grazing (so may require rebar around pvc visible above ground)

Appendix IV - Modified Hink and Ohmart categories, from NMRAM

The following is pages 39-41 in Muldavin et al.'s 2014 NMRAM for Montane Riverine Wetlands v 2.0 Manual (draft, not yet published)

Vegetation Vertical Structure Type Definitions for NMRAM

Multiple-Story Communities (Woodlands/Forests)



Type 1 – High Structure Forest with a well-developed understory.


Tall mature to intermediate-aged trees (>5 m [>15 feet]) with canopy covering >25% of the area of the community (polygon) and understory layer (0-5 m [0-15 feet]) covering >25% of the area of the community (polygon). Substantial foliage is in all height layers. (This type incorporates Hink and Ohmart structure types 1 and 3.) Photograph on Gila River by Y. Chauvin, 2012.



Type 2 –Low Structure Forest with little or no understory.


Tall mature to intermediate-aged trees (>5 m [>15 feet]) with canopy covering >25% of the area of the community (polygon) and understory layer (1-5 m [3-15 feet]) covering <25% of the area of the community (polygon). Majority of foliage is over 5 m (15 feet) above the ground. (This type incorporates Hink and Ohmart structure types 2 and 4.) Photograph on Diamond Creek by Y. Chauvin, 2012.

Single-story Communities (Shrublands, Herbaceous and Bare Ground)



Type 5 –Tall Shrub Stands.

Young tree and shrub layer only (15-5 m [4.5-15 feet]) covering >25% of the area of the community (polygon). Stands dominated by tall shrubs and young trees, may include herbaceous vegetation underneath the woody vegetation. Photograph on San Francisco River by Y. Chauvin, 2012.




Type 6S-Short Shrub Stands.

Short stature shrubs or very young shrubs and trees (up to 1.5 m [up to 4.5 feet]) covering >10% of the area of the community (polygon). Stands dominated by short woody vegetation, may include herbaceous vegetation underneath the woody vegetation. Photograph on Lower Pecos River by E. Lindahl, 2008.




Type 6W-Herbaceous Wetland.

Herbaceous wetland vegetation covering >10% of the area of the community (polygon). Stands dominated by obligate wetland herbaceous species. Woody species absent, or <10% cover. Photograph of *Carex nebrascensis* meadow on upper Rio Santa Barbara by Y. Chauvin, 2009.



Type 6H- Herbaceous.

Herbaceous vegetation covering >10% of the area of the community (polygon). Stands dominated by herbaceous vegetation of any type except obligate wetland species. Woody species absent or <10% cover. Photograph on Diamond Creek by Y. Chauvin, 2012.



Type 7–Sparse Vegetation/Bare Ground.

Bare ground, may include sparse woody or herbaceous vegetation, but total vegetation cover <10%. May be natural in origin (cobble bars) or anthropogenic in origin (graded or plowed earth) Photograph on Lower Gila River by Y. Chauvin,2012.