

Tijeras Creek Project 17-08

Pre-treatment Monitoring Report

2017



Prepared by

Kathryn R Mahan, Ecological Monitoring Specialist & Carmen Briones, Ecological Monitoring Technician,
New Mexico Forest and Watershed Restoration Institute
for the Greater Rio Grande Watershed Alliance



Ciudad SWCD

Contents

Acronyms and Abbreviations	3
Purpose of Report	4
Ecological Context of Bosque Restoration	4
Monitoring and Field Methods	5
Low intensity Field Methods	5
High-intensity Field Methods	6
Personnel Involved	6
Tijeras Creek Project	7
Discussion	17
References	18
Appendix I - Photopoint Table	20
Appendix II - Photos	21
Appendix III – Current monitoring methods available	25
Appendix IV - Modified Hink and Ohmart categories, from NMRAM	26

Acronyms and Abbreviations

Acronym, Abbreviation, or Term	Explanation or Definition as used by NMFWR
BBIRD plots	Breeding Biology Research and Monitoring Database, larger circular plot types
BEMP plots	Bosque Ecosystem Monitoring Program, small rectangular plot types
FSA	Farm Service Agency, a department of the USDA
GIS	Geographic Information Systems
GRGWA	Greater Rio Grande Watershed Alliance
LIDAR	Light detecting and ranging, a remote sensing technique using light to gather elevation data
NAIP	National Agriculture Imagery Program (aerial imagery)
NDVI	Normalized Difference Vegetation Index; GIS term for a band ratio of the visible red and the near infrared spectral bands and is calculated using the following formula: $(NIR - Red)/(NIR + Red)$
NHNM	Natural Heritage New Mexico
NMDGF	New Mexico Department of Game and Fish
NMED SWQB	New Mexico Environment Department Surface Water Quality Bureau
NMFWR	New Mexico Forest and Watershed Restoration Institute
NMHU	New Mexico Highlands University
NMRRAM	New Mexico Rapid Assessment Method, version 2.0
NRCS	Natural Resource Conservation Service
PC	Plot center
RGIS	Resource Geographic Information System
SWCD	Soil and Water Conservation District
TIFF	Tagged image file format
USDA	United States Department of Agriculture
USGS	United States Geological Survey
WQCC	Water Quality Control Commission
WSS	Web Soil Survey, a soils database of the NRCS

Purpose of Report

This report covers the low-intensity and high-intensity pre-treatment vegetation monitoring assessment performed on a non-native phreatophyte removal project submitted for the Tijeras Creek to the Greater Rio Grande Watershed Alliance in 2017. Following a discussion of the ecological context, and our monitoring methods, we present pertinent background, observations, and assessment results for the project.

Ecological Context of Bosque Restoration

Neither the challenges nor the importance of working in the bosque and other riparian areas in New Mexico today should be underestimated. According to the New Mexico Department of Game and Fish Conservation Division, wetlands and riparian areas comprise approximately 0.6 percent of all land in New Mexico (2012). Despite this small percentage, estimates of New Mexican vertebrate species depending on wetland and riparian habitat for their survival ranges from 55% (New Mexico Department of Game and Fish Conservation Services Division, 2012) to 80% (Audubon New Mexico, 2013). These areas also provide flood mitigation, filtration of sediment and pollutants, and water for a variety of purposes including groundwater recharge (Audubon New Mexico, 2013). In addition, native vegetation such as cottonwoods have cultural significance to many communities.

As much as these areas are disproportionately important to ecosystems and human communities, they are equally disproportionately impacted by disturbance. Anthropogenic impacts with major consequences for our riparian areas include dams, reservoirs, levees, channelization, acequias and ditches, jetty jacks, riprap and Gabion baskets, urbanization, removal of native phreatophytes, grazing by domestic livestock, excessive grazing pressure by native ungulate populations absent natural predation cycles, beaver removal, logging, mining, recreation, transportation, introduction and spread of invasive exotic species, groundwater extraction, altered fire and flood regimes drought and climate change (Committee on Riparian Zone Functioning and Strategies for Management, et al., 2002). Statewide, it is estimated that as much as 90% of New Mexico's historical riparian areas have been lost (Audubon New Mexico, 2013), and approximately 39% of our remaining perennial stream miles are impaired (New Mexico Department of Game and Fish Conservation Services Division, 2012).

New Mexico is fortunate enough to have the Middle Rio Grande Bosque, the largest remaining bosque in the Southwest (USDA USFS, 1996). However, over the past two decades, the number of fires in the bosque has been increasing. Historically, the primary disturbance regime in the bosque has been flooding, not fire, which means the system is not fire-adapted. In fact, native species like cottonwood resprout from their roots after floods and need wet soils to germinate from seed. Flooding also promotes decomposition of organic material and keeps the soil moist which reduces the likelihood of fire. Today, overbank flow is uncommon in many areas of the Rio Grande due to the heavy alteration of the channel and flow regimes (two obvious examples are the structures defining the upper and lower extent of the Middle Rio Grande: Cochiti Dam and Elephant Butte Reservoir). This has led to low fuel moisture content and high fuel loads, as well as increased human presence in the riparian area. As a result, bosque fires are more common and more severe: they kill cottonwoods and other native species, creating spaces which are filled by non-native species such as salt cedar, Russian olive, Siberian elm, and Tree-of-Heaven. We are constantly learning more about how these species can exploit and encourage a riparian fire regime, in addition to many other changes they bring to ecosystems.

Efforts geared toward the removal of these nonnative species can help to reduce fire risk, preserve native vegetation, and be part of a larger effort to restore the bosque and the watershed as a whole to a more natural and functional ecosystem. The Greater Rio Grande Watershed Alliance (GRGWA) has been working on these issues with a variety of collaborating organizations and agencies within the Rio Grande basin for several years. Since 2013, the New Mexico Forest and Watershed Restoration Institute (NMFWR) has been working with GRGWA and the Claunch-Pinto Soil and Water Conservation District (SWCD) to begin construction of a geodatabase for all of GRGWA's non-native phreatophyte removal projects as well as to perform the formal pre- and post-treatment monitoring, utilizing the field methods explained below as well as LIDAR analysis where appropriate and available.

Monitoring and Field Methods

Low intensity Field Methods

Low intensity pre-treatment vegetation monitoring was done using an adapted version of the biotic portion of the New Mexico Rapid Assessment Method (NMRAM), v 2.1, updating recommendations made in the Field Manual for Greater Rio Grande Watershed Alliance (GRGWA) Riparian Restoration Effectiveness Monitoring and the GRGWA Monitoring Plan, developed by Lightfoot & Stropki of SWCA Environmental Consultants in 2012. (For a brief overview of both low and high intensity monitoring methods used by the NMFWR on GRGWA projects, please see Appendix III.)

For those not familiar, NMRAM was developed by the New Mexico Environment Department Surface Water Quality Bureau Wetlands Program and Natural Heritage New Mexico as a "cost effective, yet consistent and meaningful tool" (Muldavin, 2011) for wetland ecological condition assessment in terms of anthropogenic disturbance as negatively correlated with quality and functionality. The portions of NMRAM we utilized are Level 2 "semi-quantitative" field measurements taken at less detail than plot level (Muldavin, 2011).

Measurements taken included relative native plant community composition, vegetation horizontal patch structure, vegetation vertical structure, native riparian tree regeneration, and invasive exotic plant species cover. The underlying method for these biotic assessments was a version of the 1984 Hink and Ohmart vertical structure classification system, modified for use in the NMRAM for Montane Riverine Wetlands version 2.0 (see Appendix IV). First, vegetation communities were mapped out by patch (polyon) according to the Hink and Ohmart system. Next, the presence of (state-listed) invasives, wetland species, and the two dominant species in each strata ("tree" >15 ft, "shrub" 4.5-15 ft, and "herbaceous" <4.5 ft) were recorded for each plant community. The native/exotic ratio in each of the patches was scored and weighted based on the percent of the project area each patch comprised. These scores were then combined with the additional biotic metrics of vertical and horizontal diversity, native tree regeneration, and overall (listed) invasive presence. The NMRAM rating system is based, on all levels, on a scale of 1 to 4, where 4 is considered excellent condition, 3 good, 2 fair, and 1 poor.


We also assessed soil surface condition, which is a metric typically included in the abiotic section of the NMRAM, as well as the presence of surface fuels, which is not part of the NMRAM. Unlike the other 6 metrics we used, surface fuels were recorded on a rating scale from 0 to 1.0 where 1.0 is a continuous fuel matrix.

Photopoints were established to capture images where vegetation shifts were observed and/or at representative locations throughout the site. Waypoints were marked with a Garmin GPS unit and named sequentially by site. Photos were taken facing north, east, south and west at each point. Information about the photopoints was collected according to the methods laid out in David Lightfoot's Forest Thinning Project Repeat Photo Points for Restoration Effectiveness Monitoring (David Lightfoot, 2014).

Prior to entering the field, we created a map with the project boundaries as provided by GRGWA. We combined these polygons with recent aerial imagery and identified relevant roads and other landscape features. Once on the ground, the vegetation community polygons (as determined by the modified Hink and Ohmart classification system) were hand-drawn onto this map and served as the basis for other biotic metric assessments. Upon return to the office, this polygon map and the photopoints were digitized by the monitoring technician and/or specialist.

High-intensity Field Methods

High-intensity monitoring was also done, in part, on this site. We used an adapted Bosque Ecosystem Monitoring Program (BEMP) style plot. These are 16 x 98-foot rectangles, placed approximately parallel to the river. Within these plots, we measure canopy and species, and vegetation and ground cover. We also used Brown's transects to measure surface fuels.



Personnel Involved

2016 New Mexico Forest and Watershed Restoration Institute Monitoring Team:

- Kathryn R Mahan, Ecological Monitoring Specialist
- Ernesto Sandoval, Ecological Monitoring Technician

2016 New Mexico Forest and Watershed Restoration Institute GIS Team:

- Daniel Hernandez, Ecological Monitoring Technician

Other persons contacted:

- Fred Rossbach, Field Coordinator, Greater Rio Grande Watershed Alliance
- Marcos Valdez, East Rio Arriba Soil and Watershed Conservation District
- Jim Brooks, Soilutions
- Sarah Hurteau, TNC
- Kali Bronson, Bernalillo County

- Laurie Wearne, Talking Talons Youth Leadership
- Michael Young, Adaptive Terrain Systems

Tijeras Creek Project

Project 17-08 Tijeras Creek is 12.06 acres located North of A. Montoya Elementary School, South of Highway 333 and Los Vecinos Community Center in the Village of Tijeras, in the City of Albuquerque, New Mexico. Summer 2017 an Ecological monitoring took place at the Tijeras Creek by different members like the Arid Land Innovation, The Nature Conservancy, Ecotone Landscape Planning and Adaptive Terrain systems. Monitoring was conducted by staff members from Talking Talons Leadership, and several monitoring methods and workshops were addressed in order to train and approach the restoration goals for the site. One of the workshops covered ID of non-native species and their management. The monitoring design included reduction of invasive species, increase of native plant community, and improvement of hydrologic functions. Data was collected on an adequate sample size plots, a total of 14 transects were fixed and crew members recorded, measured and took note on understory cover of herbaceous species. Based on the results of the assessment the monitor report recommends maintenance of invasive species, photo records and use of game camera for wildlife use on restored area. Future monitoring component suggests to go out and recollect data on the measurements of vegetation to evaluate the effectiveness of the restoration goal. (Talking Talons Youth Leadership, 2017). For more details and full coverage of the Tijeras Creek Ecological Monitoring report you can contact the Talons Youth Leadership at Post Office Box 8 Cedar Crest, NM 87008, or (505) 281-1133. The link to their website is <http://www.talkingtalons.org/>

Pre-treatment monitoring was conducted on October 24, 2017 as part of a restoration project targeting the ecological functions, like removing non-native species, stabilizing streamside banks, and replanting native riparian plants. Site accesses is via Public School Road (paved) and from Los Vecinos Center (by foot). The project was proposed in 2017, sponsors are Ciudad Soil and Water Conservation District (CSWCD) and the Tijeras Creek Watershed Collaborative. Degraded conditions are contributed by several noxious weed patches and colonization of Siberian elm trees. Desired site condition would be to have a riparian zone of native species with overstory canopy cover greater than 40%, and 10% of canopy cover of Siberian elm and 0% of ground cover representing invasive species. Native species are Rio Grande cottonwood, juniper spp, smooth brome and chamisa. Invasive species found at this site are Siberian elm, Dalmatian toadflax, salt cedar and cheatgrass. Treatment type proposed is hand crew extraction, by cutting stumps and chipping would also be ok as it be utilized for woody material cover. For this project it has been considered to leave larger elms for now as they provide shade. Large leave trees will be removed with a follow up project. Stream bed modifications can be incorporated into the non-native treatments. Talking Talons will continue monitoring combined with proposed erosion control.


The average precipitation for Albuquerque is 9.39 inches annually. The annual high temperature average is 68.8° F, the average low is 45.4° F and average temperature is 57.1°F (U.S. Climate Data, 2017). According to the NRCS Web Soil Survey, the project area is comprised of 77.3% Manzano loam and 22.7% La Fonda loam. Rangeland ecological sites within this project include R070CY109NM Loamy, R035XG121NM Shallow Sandstone and R042XA054NM Deep Sand (USDA NRCS, 2016).

The Loamy ecological site typical plant community is a mixed grassland of warm and cool season, mid- and short perennial grasses. Woody species occupy a minor portion of this community, as do forbs. During periods of abundant spring and fall moisture, a large variety of forbs can be found. Piñon and oneseed juniper can occur in some portions of this site (USDA NRCS n.d., n.d.).

The Shallow Sandstone ecological site type is dominated by grasses such as sideoats grama, blue grama, little bluestem, Indian ricegrass, New Mexico feathergrass, and galleta. Shrubs found in this site type include Bigelow sagebrush and fourwing saltbush. Other common shrubs include sand sagebrush, rubber rabbitbrush, winterfat, and mountain mahogany. In its reference condition, piñon and juniper are scattered across this site type, however, grasses are dominant with fairly uniform cover and few large bare areas present. Scattered shrubs and trees may comprise a canopy cover averaging 10%. Evidence of erosion such as pedestalling of grasses, rills and gullies is infrequent. Sideoats grama, little bluestem, many cool-season grasses, mountain mahogany, and winterfat typically decrease in response to overgrazing resulting in a blue-grama/galleta community or even a piñon-juniper dominated community when overgrazing is combined with fire suppression/lack of fine fuels and mild summers paired with wet winters which favor juniper establishment (USDA NRCS n.d.).

The Deep Sand ecological site type is mainly grassland and quite an amount of shrubs. The grasslands consists of a mixture of short-, mid-, and tall grasses. Annual grasses and forbs occur in relatively large amounts. Plant community include: six-weeks grama, sand muhly, blue grama, foxtail barley, bottlebrush squirreltail, tumblegrass and threeawn spp. Other forbs include: tansymustard, stickleaf, globemallow, silverleaf nightshade, locoweed, woolly grounseel, and indian paintbrush. When the plant community deteriorates, there is an increase of woody and succulent plants. Mesquite and juniper may overtake in the site. In severe conditions of worsening of plant community, there will be active soil erosion resulting in bared sand dunes (USDA NRCS, n. d.).

17-08 Tijeras Creek



1 inch = 755 feet

Project 17-08 Tijeras Creek, is located in Bernalillo County in the town of Tijeras, New Mexico. N.M. HWY 333 borders the northern part of the project area and Public School road outlines the southern boundary. The project is considered an initial treatment with Siberian Elm, salt cedar and juniper as the target species.



Map by Daniel Hernandez

Figure 1. Project 17-08 in geographic context.




Figure 2. 17-08 Tijeras Creek project outline.

Table 1. NMRAM Scores for 17-08.

Metric 17-08, 24 Oct 17	Score
Relative Native Plant Community Composition	1
Vegetation Horizontal Patch Structure	4
Vegetation Vertical Structure	3
Native Riparian Tree Regeneration	3
Exotic Invasive Plant Species Cover	1
Project Biotic Score (based on above ratings)	2.2
Project Biotic Rating	C/ Fair
Soil Surface Condition	3
Surface Fuels	0.6

The lowest scores for this project came in the Relative Native Plant Community Composition and Exotic Invasive Plant Species Cover metrics, due to the high percentage of invasive plants. The project scored best in the vegetation horizontal metrics, because there are several different plant communities distributed across the landscape. Vegetation polygons are represented by structure type in the map, Figure 3. This site scored a 2.2 out of 4 overall, which is a “C” or “Fair” biotic rating.

In this site a map was created (Figure 4) to showcase the center of the community and the heart of Tijeras which is Los Vecinos Area. The map shows a detailed plan of trails that connect to facilities and areas with an easy access amending to the requests of the community in the area.

17-08 Tijeras Creek




Figure 3. 17-08 Tijeras Creek project vegetation polygons.




Figure 4. Los Vecinos Area Plan for 17-08.

Table 2. 17-08 Average surface fuels from 2 transects on plot.

Fuel	Average tons/acre
1-hr	0.35
10-hr	0
100-hr	0.21
1000-hr	0.00
All woody fuels	0.56
Fuel	Avg depth (inches)
Duff	0
Litter	0.11
Total	0.11




Figure 5. Percent Vegetative Cover for plot on 17-08.




Figure 6. Percent Vegetative Cover for plot on 17-08.




Figure 7. Percent ground cover for plot on 17-08.




Figure 8. Percent ground cover for plot on 17-08.

Discussion

We would like to clarify that we are adapting these NMRAM metrics for our own purposes. That is, we are using them both inside and outside their intended site ranges, including on larger sites (NMRAM is designed to handle a site around 100 x 200 meters), sites further from the river (NMRAM is currently in use primarily for assessing riverine wetlands), and sites defined by exotic vegetation presence rather than hydrologic boundaries and upland vegetation indicators/apparent wetland extent. Site delineation and size is likely to be variable for a number of other reasons, including landowner participation, available funds, proposals received from contractors, etc – many of which cannot be directly correlated to site disturbance or ecological function. For this reason, we do not use the entire NMRAM assessment, or place confidence in the weighted score roll-ups that are typically part of an NMRAM report. Should one be interested, rationale for the weighting in the NMRAM score roll-up can be found in the yet-to-be-published field manual for version 2.1. For more information, contact Maryann McGraw of the NMED or NMFWR.

While we provide a biotic site score and rating for your reference, we recommend comparisons be done with individual metrics from pre-treatment and post-treatment assessment from the same site, rather than across multiple sites. Also of note is that statistical analysis is not appropriate for NMRAM, or other low intensity, rapid field methods.

Please note that should the project area change significantly from what was originally proposed and monitored, all metrics will lose some amount of confidence on comparison as it is impractical to re-examine the original site assessment scores using new boundaries. This is an issue of concern of which GRGWA should be aware. We recommend that GRGWA attempt to minimize alterations in project boundaries once pre-treatment monitoring data has been approved for collection. Another, somewhat alternative, recommendation is that the initial monitoring regime include high-intensity modified BEMP-type plots which could be repeated in their exact initial locations, allowing collection of comparable data regardless of boundary change. We recognize that this is not always practical: boundaries change for a number of reasons and time and cost constraints can necessitate the sole use of a rapid assessment method for monitoring. We have reason to hope our outlined assessment method will still be a satisfactory indicator for site function improvement or degradation primarily because metrics in rapid assessment methods such as this are set up to have relatively low sensitivities (i.e. for a change to be reflected in the metrics, either positive or negative, disturbance on site has to be significantly altered).

From here on out, the goal of the GRGWA/ NMFWR is that all sites will be revisited for post-treatment monitoring in 5-year intervals. It is our intention and expectation that the data collected in these intervals will reflect any significant changes in disturbance and ecological function of the site.

References

- Audubon New Mexico. (2013). *Water Matters: Water for New Mexico Rivers*. Albuquerque, New Mexico: Utton Transboundary Resources Center.
- Brown, J. K. (1974). *Handbook for Inventorying Downed Woody Material*, USDA Forest Service General Technical report INT-16. *Handbook for Inventorying Downed Woody Material*. Ogden, Utah: USDA Forest Service Intermountain Forest and Range Experiment Station.
- Bureau of Land Management. (2006). *Grazing Management Processes and Strategies for Riparian-Wetland Areas, TR 1737-20*.
- Claunch-Pinto Soil and Water Conservation District on behalf of the Greater Rio Grande Watershed Alliance. (2015). *Request for Proposals for Greater Rio Grande Watershed Alliance Riparian Restoration Projects*. Mountainair, NM: Claunch-Pinto Soil and Water Conservation District.
- Committee on Riparian Zone Functioning and Strategies for Management, et al. (2002). *Riparian Areas: Functions and Strategies for Management*. Washington, D.C.: National Academy Press.
- David Lightfoot. (2014). *Forest Thinning Project Repeat Photo Points for Restoration Effectiveness Monitoring*. Albuquerque: SWCA Environmental Consultants.
- Lightfoot, D. &. (2012). *Greater Rio Grande Watershed Alliance Riparian Restoration Effectiveness Monitoring Plan*. Albuquerque, NM: SWCA Environmental Consultants.
- Lightfoot, David & Stropki, C. (2012). *Field Manual for Greater Rio Grande Watershed Alliance Riparian Restoration Effectiveness Monitoring*. Albuquerque, NM: SWCA Environmental Consultants.
- Lizarazo, I., & Elsner, P. (2009). Fuzzy segmentation for object-based image classification. *International Journal of Remote Sensing*, 30.
- Muldavin, E. B. (2011). *New Mexico Rapid Assessment Method: Montaine Riverine Wetlands*. Version 1.1. Final report to the New Mexico Environment Department, Surface Water Quality Bureau. 90 pp. and appendices.
- New Mexico Department of Game and Fish Conservation Services Division. (2012). *Bridge and Road Construction/Reconstruction Guidelines for Wetland and Riparian Areas*.
- Talking Talons Youth Leadership. (2017). *Tijeras Creek Ecological Monitoring Report*. Albuquerque.
- U.S. Climate Data. (2017). *Climate New Mexico*. Retrieved from U.S. Climate Data: <http://www.usclimatedata.com/climate/new-mexico/united-states/3201>
- USDA NRCS. (2016, 8 10). *Web soil Survey*. Retrieved from <https://websoilsurvey.sc.egov.usda.gov/App/HomePage.htm>
- USDA NRCS. (n. d.). *Ecological Site Description for Deep Sand*. Retrieved February 21, 2018, from <https://esis.sc.egov.usda.gov/ESDReport/fsReport.aspx?approved=yes&rptLevel=all&id=R042XA054NM>
- USDA NRCS n.d. (n.d.). *Ecological Site Description Shallow Sandstone R035XG121NM*.

USDA NRCS n.d. (n.d.). *Ecological site characteristics loamy R070CY109NM*. Retrieved from Ecological site information system:
<https://esis.sc.egov.usda.gov/ESDReport/fsReport.aspx?approved=yes&repType=regular&id=R070CY109NM>

USDA USFS. (1996, September). *Ecology, Diversity, and Sustainability of the Middle Rio Grande Basin*, RM-GTR-268. (D. M. Finch, & J. A. Tainter, Eds.) Fort Collins, Colorado.

Appendix I - Photopoint Table

Name	Latitude	Longitude
17.08_1_N	35.07794	-106.39292
17.08_1_E	35.07794	-106.39292
17.08_1_S	35.07794	-106.39292
17.08_1_W	35.07794	-106.39292
17.08_2_N	35.07888	-106.39555
17.08_2_E	35.07888	-106.39555
17.08_2_S	35.07888	-106.39555
17.08_2_W	35.07888	-106.39555
17.08_PC_NESW	35.07862	-106.39505

Appendix II - Photos



Tijeras Creek 17.08_1_N
View facing north inside of
polygon 5. Taken from the
bridge.

17.08_1_E View facing
east inside of polygon 5
and 3. Taken from the
bridge.



17.08_1_S View facing south
inside of polygon 3. Taken
from the bridge.

17.08_1_W View facing
west inside of polygon 3 and
5. Taken from the bridge.



17.08_2_N View facing
north inside of polygon 9,
8 and 5. Foreground
landmark are elms.

17.08_2_E. View facing
east inside of polygon 9
and 5.





17.08_2_S View facing south inside of polygon 9 and 2. Foreground landmarks are elm sprouts.


17.08_2_W View facing west inside of polygon 9. Foreground landmark is a razor wire @ 260° approx. 150ft away.



17-08 Plot Photos




N from PC



E from PC



S from PC



W from PC



Plot Center

Appendix III – Current monitoring methods available

Low-intensity methods

- Where: happens on all sites with GRGWA projects
- Method name: NMRAM (New Mexico Rapid Assessment Method v 2.1)
- Time required: 3 hours – half day/ site
- Repeat: done once pre-treatment and in 4-5 year intervals post-treatment
- Basics: mapping vegetation communities (by vertical and horizontal structure), recording dominant vegetation in each strata (trees, shrubs, herbaceous), assessing fuel load, noting soil surface condition and native/exotic ratio at all vegetation levels, photo points
- Any on-site impacts or materials: none

High-intensity methods

- Where: happens on select sites, in addition to low-intensity monitoring

Submethod name 1: BBIRD or BEMP vegetation plots (depends on treatment area size)

- Time required: approx. 2 hours/site
- Repeat: both pre-treatment and in 4-5 yr intervals post-treatment
- Basics: larger plots and transects documenting vegetation, photo points
- On-site impacts or materials: rebar and cap

Submethod name 2: Brown's transects

- Time required: 1-1.5 hours/site
- Repeat: both pre-treatment and in 4-5 yr intervals post-treatment
- Basics: transects to calculate fuel loading and fire behavior, photo points
- On-site impacts or materials: rebar and cap

Submethod name 3: BEMP-adapted Groundwater Well Monitoring

- Time required:
 - Initial installation: 1-2 hours/ well (ideally 2+ wells/site)
 - Repeat: maintenance as needed, should be minimal
 - Data offloading: 10-20 minutes/well
 - Repeat: at least annually (this is when we anticipate datalogger will be full and batteries will need to be changed)
- Basics: install a well with a sensor which records groundwater level and temperature once an hour year round; this will reflect changes due to seasonal variation, vegetation growth, irrigation, etc.
- On-site impacts or materials: shallow monitoring well (consists of capped PVC pipe extending into the ground about 3 feet below the water table and above ground approx. 2 feet (can be painted earth tones); well contains a datalogger (pressure transducer) suspended on a cable into the water); well should be protected from cattle grazing (so may require rebar around pvc visible above ground)

Appendix IV - Modified Hink and Ohmart categories, from NMRAM

The following is pages 39-41 in Muldavin et al.'s 2014 NMRAM for Montane Riverine Wetlands v 2.0 Manual (draft, not yet published)

Vegetation Vertical Structure Type Definitions for NMRAM

Multiple-Story Communities (Woodlands/Forests)



Type 1 – High Structure Forest with a well-developed understory.

Tall mature to intermediate-aged trees (>5 m [>15 feet]) with canopy covering >25% of the area of the community (polygon) and understory layer (0-5 m [0-15 feet]) covering >25% of the area of the community (polygon). Substantial foliage is in all height layers. (This type incorporates Hink and Ohmart structure types 1 and 3.) Photograph on Gila River by Y. Chauvin, 2012.



Type 2 –Low Structure Forest with little or no understory.


Tall mature to intermediate-aged trees (>5 m [>15 feet]) with canopy covering >25% of the area of the community (polygon) and understory layer (1-5 m [3-15 feet]) covering <25% of the area of the community (polygon). Majority of foliage is over 5 m (15 feet) above the ground. (This type incorporates Hink and Ohmart structure types 2 and 4.) Photograph on Diamond Creek by Y. Chauvin, 2012.

Single-story Communities (Shrublands, Herbaceous and Bare Ground)



Type 5 –Tall Shrub Stands.


Young tree and shrub layer only (15-5 m [4.5-15 feet]) covering >25% of the area of the community (polygon). Stands dominated by tall shrubs and young trees, may include herbaceous vegetation underneath the woody vegetation. Photograph on San Francisco River by Y. Chauvin, 2012.

**Type 6S-Short Shrub Stands.**


Short stature shrubs or very young shrubs and trees (up to 1.5 m [up to 4.5 feet]) covering >10% of the area of the community (polygon). Stands dominated by short woody vegetation, may include herbaceous vegetation underneath the woody vegetation. Photograph on Lower Pecos River by E. Lindahl, 2008.

**Type 6W-Herbaceous Wetland.**

Herbaceous wetland vegetation covering >10% of the area of the community (polygon). Stands dominated by obligate wetland herbaceous species. Woody species absent, or <10% cover. Photograph of *Carex nebrascensis* meadow on upper Rio Santa Barbara by Y. Chauvin, 2009.

**Type 6H- Herbaceous.**

Herbaceous vegetation covering >10% of the area of the community (polygon). Stands dominated by herbaceous vegetation of any type except obligate wetland species. Woody species absent or <10% cover. Photograph on Diamond Creek by Y. Chauvin, 2012.



Type 7–Sparse Vegetation/Bare Ground.

Bare ground, may include sparse woody or herbaceous vegetation, but total vegetation cover <10%. May be natural in origin (cobble bars) or anthropogenic in origin (graded or plowed earth) Photograph on Lower Gila River by Y. Chauvin, 2012.